

Efficacy of Thirteen Species of Wild Flora as Soil Amendments in the Control of the Root-Knot Nematode, *Meloidogyne javanica* on Common Bean in Saudi Arabia

Amr A. El-Sherbiny¹; and Fahad A. Al-Yahya²

ABSTRACT

Powders of the aerial parts of thirteen Saudi wild plant species namely: *Achillea fragrantissima*, *Anastatica hierochuntica*, *Brassica sinaica*, *Calotropis procera*, *Cleome rupicola*, *Ducrosia anethifolia*, *Fagonia bruguieri*, *Lactuca serriola*, *Pulicaria crispa*, *Reseda muricata*, *Salsola imbricata*, *Trichodesma africanum* and *Zygophyllum migahidii* were screened for their nematicidal activity towards the root-knot nematode (*Meloidogyne javanica*) infecting common bean cv. Strike in a pot experiment under greenhouse conditions ($35\pm5^{\circ}\text{C}$). Plant powders were applied as amendments of nematode-infested soil at 0.3% (w/w), comparing to a treatment with the nematicide, Carbofuran 10%G (0.1g/pot), 7 days prior to planting. Nematode-free soil and nematode-infested soil were served as checks. All studied plant powders significantly reduced ($P\leq0.05$) numbers of root galls, nematode egg masses, final nematode egg populations and reproduction factor on common beans, as compared to nematode check. Among them, *C. rupicola*, *S. imbricata*, *T. africanum* and *Z. migahidii* provided the maximum reductions of all disease parameters ranged from 82.0-96.4%, whereas *F. bruguieri* gave the minimum ones (33.9-40.4%). Meanwhile, application of Carbofuran 10% resulted in 88.0-90.8% reduction of all disease parameters. On the other hand, common bean growth of the most treatments significantly increased ($P\leq0.05$), as compared to nematode check plants, and mostly to the nematicide-treated and healthy ones. Percentages of increase were ranged from 80.1 - 188.3% in shoots and 106.8 - 278.8% in roots. On the contrary, soil treatments with the powder of *A. hierochuntica* and *Z. migahidii*, showed little phytotoxicity symptoms on common beans. Powders of *C. rupicola*, *S. imbricata*, *T. africanum* and *Z. migahidii* achieved relative nematicidal efficacy ranged from 66.5 - 269.1% of Carbofuran 10%.

The current results recorded new wild plants in Saudi Arabia with a potential nematicidal activity against the root-knot nematode. They may be considering a safe source of new alternative nematicides.

INTRODUCTION

The root-knot nematode (*Meloidogyne* spp.) is a real threat and damaging pest of vegetable crops and many economic plants growing in Saudi Arabia and

worldwide (Al-Hazmi *et al.*, 1995). Its management has been relied mainly on chemical nematicides, but due to the environment pollution caused as a result of repeated application of chemical pesticides, several attempts were devoted towards recent approaches for safe nematode management. Pesticidal chemicals of plant origin have received more attention by many researchers because of they are less concentrated than synthetic ones, biodegrade rapidly and they can be derived from renewable natural resources (Quarles, 1992).

In Saudi Arabia, little trials were carried out to study the nematicidal activity of indigenous wild flora. Study of Al-Rajhi *et al.*, (1997) revealed that *Juniperus polycarpos* and *Rhazya stricta* (plants wildly grown near Riyadh) have *in vitro* nematicidal properties towards *Meloidogyne javanica*. Similarly, Al-Yahya *et al.*, (2005) gave a short report on the nematicidal activity of 14 different species of Saudi wild plants against *M. javanica* under laboratory conditions.

Fortunately, Saudi Arabia is gifted by a great diversity of natural vegetations and wild plants in its deserts. Among the common plants wildly growing in Saudi deserts are: *Achillea fragrantissima*, *Anastatica hierochuntica*, *Brassica sinaica*, *Calotropis procera*, *Cleome rupicola*, *Ducrosia anethifolia*, *Lactuca serriola*, *Pulicaria crispa*, *Reseda muricata*, *Salsola imbricata*, *Trichodesma africanum*, *Fagonia bruguieri*, and *Zygophyllum migahidii* (Chaudhary, 1999; Chaudhary and Al-Jowaid, 1999).

Indeed, most of the above mentioned wild plants have therapeutic properties and traditionally used in the folk medicine to cure and/or reduce symptoms of various disorders including infectious diseases among Bedouins (Salamah *et al.*, 1989; Mossa *et al.*, 1991; Tanira *et al.*, 1994; Rahman *et al.*, 2004; Bogdadi *et al.*, 2007; Abdel-Sattar *et al.*, 2010; Ahmed *et al.*, 2010 and Maghraby *et al.*, 2010).

Nothing was found in the literature regarding the nematicidal activity of the most studied wild plants, except for *A. fragrantissima* (Oka *et al.*, 2000) and *C. procera* (Nandal and Bhatti, 1990; Rao *et al.*, 1996).

¹Nematology Research Dept., Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt
E-mail: amr_elsherbiny_1968@yahoo.com

²Plant Protection Dept., College of Food and Agricultural Sciences, King Saud University, Riyadh 11451, P.O. 2460, Saudi Arabia
Received December 1, 2010, Accepted December 29, 2010.

However, some available informations concerning antimicrobial and/or toxic activities of these plants were reported. *A. hierochuntica* showed strong antibacterial, anticandidal and antifungal activities against certain species of Gram positive and Gram negative bacteria, *Candida* spp., and *Aspergillus* spp. (Mohamed *et al.*, 2000; Mohamed *et al.*, 2009).

Cleome rupicola (syn. *Cleome arabica*) is a wild plant with a distinct foetid smell. It has an effective antibacterial activity against *Klebsiella pneumoniae* and *Staphylococcus aureus* (Salamah *et al.*, 1989). Moreover, *C. arabica* and other *Cleome* species wildly growing in Saudi Arabia have antimicrobial, anthelmintic, antiseptic and toxic activities (Rahman *et al.*, 2004). Similarly, *D. anethifolia* have a strong aromatic odour and it has antimycobacterial, antifungal, antibacterial and anticandidal activities (Janssen *et al.*, 1984; Stavri *et al.*, 2003; Pirbalouti, 2009).

Crude methanolic extract of *L. serriola* posses antiplasmodial, antitrypanosomal and antilishmanial activities (Abdel-Sattar, *et al.*, 2010).

Pulicaria crispa (syn. *Francoeuria crispa*) is commonly named as Gethgath in Saudi Arabia. It has multiple antimicrobial and pesticidal activities including: antiplasmodial (Sathiyamoorthy *et al.*, 1999), insecticidal (Al-Doghairi and Elhag 2003), strong antibacterial and anticandidal (Bogdadi *et al.*, 2007) and anthelmintic activities (Maghraby *et al.*, 2010).

Extracts and the essential oil of *T. africanum* showed strong antibacterial, antifungal, anticandidal (Abd El-Moaty, 2009), antiplasmodial, antitrypanosomal and antilishmanial activities (Abdel-Sattar, *et al.*, 2010).

Beside the nematicidal activity of *A. fragrantissima* (Oka *et al.*, 2000), it has also antiplasmodial (Sathiyamoorthy *et al.*, 1999), antibacterial, antifungal and anticandidal activities (Al-Gaby and Allam, 2000).

Herein, we studied the nematicidal efficacy of thirteen species of Saudi wild plants against the root-knot nematode, *Meloidogyne javanica* infecting common bean (*Phaseolus vulgaris* L.) cv. Strike, comparing to the chemical nematicide Carbofuran 10%G under greenhouse conditions.

MATERIALS AND METHODS

Plant materials:

Fresh aerial parts of eleven species of Saudi wild plants belonging to nine families namely: *Brassica sinaica* (Brassicaceae), *Calotropis procera* (Asclepiadaceae), *Cleome rupicola* (Capparaceae), *Ducrosia anethifolia* (Apiaceae), *Fagonia bruguieri*, *Zygophyllum migahidii* (Zygophyllaceae), *Lactuca serriola*, *Pulicaria crispa* (Asteraceae), *Reseda muricata* (Resedaceae), *Salsola imbricata*

(Chenopodiaceae) and *Trichodesma africanum* (Boraginaceae) were collected from different localities of western Riyadh deserts during the period from January-March, 2005. Dried whole plants of *Anastatica hierochuntica* (Brassicaceae) and aerial parts of *Achillea fragrantissima* (Asteraceae) were purchased from an herbal shop at Riyadh city. The studied plants were botanically identified according to descriptions given by Chaudhary, (1999) and Chaudhary and Al-Jowaid (1999), left to air drying on a clean bench at lab temperature for 10-15 days till complete dryness, well-ground in a coffee grinder and sieved through a 100 mesh stainless steel sieve (150 μ pore aperture) to obtain the fine powders.

Nematode culture and inoculum:

The root-knot nematode, *Meloidogyne javanica* (Treub) Chitwood, originally isolated from severely galled eggplants grown in Riyadh, was cultured on tomatoes (*Lycopersicon esculentum* Mill.) cv. Marmande in the greenhouse for 60 days. For soil infestation, nematode eggs were extracted from galled tomato roots using 0.5% sodium hypochlorite (Hussey and Barker, 1973).

Experiment procedures:

Clean plastic pots of 14cm diam (ca. 1300g soil) were filled with a steam sterilized soil mixture of sand, silt, and ground peat moss (4:2:1, v:v:v). Soil mixture was weighted before placing in the pots. Powders of the studied wild plants at 0.3% w/w (4g/pot) were mixed thoroughly with the potted soils. For comparison, granules of the nematicide Carbofuran 10% (0.1g/pot) were mixed with a potted soil in a separate treatment. Each treated soil was thoroughly infested with 20ml of nematode eggs suspension containing 4500 eggs. Nematode-free soil and soil infested with nematode only were kept as checks. All treated pots were watered upto the field capacity for 7 days in order to decompose plant powders (Dahiya, 2003). Treatments were replicated five times and arranged in a complete randomized design on a clean bench in the greenhouse (air temperature 35±5°C). Following the 7 days of decomposition, each pot was planted with three common bean seeds, and seedlings were thinned to one/pot, three days after emergence and received their needs of water and fertilization. Fifty days after planting, plants were carefully uprooted and roots were gently washed with running tap water, stained for 15 minutes in an aqueous solution of Phloxine B (0.15g/L. water), then rewashed with running tap water to remove the residuals of stain and emphasize nematode egg masses for counting (Holbrook *et al.*, 1983). Numbers of root galls and nematode egg masses were counted using a hand counter. Moreover, final nematode egg populations (Pf)

in the egg masses in common bean roots were extracted by the sodium hypochlorite method (Hussey and Barker, 1973) and counted, then nematode reproduction factor (Rf) was calculated (Oostenbrink, 1966). Fresh shoots and roots of common beans were weighted prior to root staining.

Statistical analysis:

Nematode disease parameters (including numbers of root galls, nematode egg masses, nematode Pf and Rf values) and plant growth parameters were subjected to the analysis of variance (ANOVA) using SAS software program (SAS, 1997). Means of all treatments were compared with Fisher's protected LSD at 5% level of probability.

RESULTS AND DISCUSSION

All tested plant powders achieved good control of *M. javanica* on common beans, as compared to nematode check plants. The percentages of nematode reduction ranged from (33.9 - 40.4%) for *F. bruguieri* to (88.2 - 96.4%) for *T. africanum*. Also, Carbofuran 10%G provided considerable nematode control ranged from 88.0 - 90.8% (Table 1).

Numbers of final egg populations (Pf) and reproduction factor (Rf values) of *M. javanica* greatly ($P \leq 0.05$) suppressed by soil treatment with all tested plant powders at 0.3% w/w (Table 1). Generally, these findings are similar to the results of Oka, *et al.*, (2001). They found that leaf powder of *Inula viscosa* (a wild plant from Palestine) when mixed to *M. javanica*-infested soil at a low concentration (0.1% w/w) greatly reduced the number of second-stage juveniles of the nematode in a sandy soil.

Soil treatment with powders of *A. fragrantissima* significantly reduced (60.6 – 68.6%) nematode infection on common beans. Nematicidal effects of *A. fragrantissima* essential oil on egg hatching and mobility of second stage juveniles of *M. javanica* under laboratory conditions were previously confirmed by Oka *et al.*, (2000). Moreover, powders of *B. sinaica* controlled *M. javanica* on common beans with acceptable reduction (48.5 - 61.4%). This result is agreed with that given by Ibrahim *et al.*, (1994), who found that soil amendment with powders of cauliflower leaves (*Brassica oleracea* var. *botrytis*) at 2% (w/w) gave a considerable reduction (72-80%) of *M. incognita* on corn. Also, soil treatment with chopped broccoli leaves (*Brassica oleracea* var. *botrytis*) at 2% (w/w) significantly reduced *M. incognita* and *M. javanica* infections on melon plants (Ploeg and Stapleton, 2001).

C. procera provided a satisfied reduction of *M. javanica* (61.7 - 67.6%) on common beans. This finding confirmed the earlier reports on the nematicidal

potential of this plant in the management of root-knot nematodes on eggplant (Nandal and Bhatti, 1990) and tomato (Rao *et al.*, 1996).

Phytochemical constituents of the studied wild plants are summarized in Table 3. Chitwood, (2002) documented that naturally occurring phytochemicals antagonistic towards plant-parasitic nematodes found in higher plants are including: polythienyls; isothiocyanates and glucosinolates (from Brassicaceae); cyanogenic glycosides; polyacetylenes (from Asteraceae); alkaloids; fatty acids and their derivatives; essential oils; terpenoids; sesquiterpenoids; diterpenoids; quassinoids; steroids; triterpenoids; and phenolics. Therefore, it was suggested that nematicidal activity of the studies plants against *M. javanica* may be attributed to one or more of the listed phytochemical constituents (Table 3).

Growth parameters of common bean of the most treatments were significantly ($P \leq 0.05$) increased, as compared to nematode check plants and mostly to the nematicide-treated and healthy ones. Percentages of increase were ranged from 80.1 - 188.3% in shoots and 106.8 - 278.8% in roots (Table 2). On the contrary, powders of *A. hierochuntica* and *Z. migahidii* showed little phytotoxicity symptoms. Ibrahim and Ibrahim (2000) reported that leaf powders of *Eucalyptus* sp. and *Psidium guagava* caused phytotoxicity on common beans cv. Giza 3. Although, soil application with powders of *F. bruguieri* gave the minimum nematode reduction (Table 1), it greatly enhanced plant growth (Table 2). Therefore, it is suggested that increase in plant growth may be attributed to nematode reduction caused by one or more of the phytochemical constituents (Table 3) and/or nutrients released during the decomposition of plant powders as organic soil amendments. It seemed that most studied plant powders improve soil fertility. Nematode suppression caused by organic soil amendments have a lot of possibilities including release of pre-excising nematicidal compounds in soil amendments, generation of nematicidal compounds such as ammonia and fatty acids during decomposition, increase in plant tolerance to nematode infection, and changes in soil physiology and structure that are unsuitable for nematode behavior (Oka, 2010).

Based on the relative efficacy of all studied plant powders to the nematicide, Carbofuran 10%G (Table 1), it is clear that *T. africanum* was the best plant in suppressing nematode infection on common beans with respectable nematicidal efficacy (101.1–269.1%), followed by *C. rupicola*, *S. imbricata* and *Z. migahidii* with nematicidal efficacies ranged from 66.5–102.4%. On the other hand, *P. crispa* and *R. muricata* gave a nematicidal efficacy slightly less than half (33.8–44.4%) of the nematicide.

Table2. Effect of powders of thirteen species of Saudi wild plants as soil amendments at 0.3%(w/w) and Carbofuran 10%G (0.1g/pot) on growth of common bean cv. Strike, 50 days after planting in *Mj*-infested soil under greenhouse conditions (35±5°C)

Treatment	Fresh weight (g)			
	Shoot	Increase (%)*	Root	Increase (%)
Nematode check (<i>Mj</i> only)	2.06 g	—	1.32 i	—
Healthy plants (nematode-free)	3.08 ef	49.5	2.25 gh	70.5
Carbofuran 10%G + <i>Mj</i>	3.06 ef	48.5	2.00 h	51.5
<i>Achillea fragrantissima</i> + <i>Mj</i>	4.56 bc	121.4	3.13 ef	137.1
<i>Anastatica hierochuntica</i> + <i>Mj</i>	2.52 fg	22.3	2.19 gh	65.9
<i>Brassica sinaica</i> + <i>Mj</i>	4.87 bc	136.4	4.09 bcd	209.8
<i>Calotropis procera</i> + <i>Mj</i>	5.00 bc	142.7	4.61 abc	249.2
<i>Cleome rupicola</i> + <i>Mj</i>	5.33 ab	158.7	4.01 cd	203.8
<i>Ducrosia anethifolia</i> + <i>Mj</i>	4.49 cd	118.0	3.55 de	168.9
<i>Fagonia bruguieri</i> + <i>Mj</i>	5.94 a	188.3	4.73 ab	258.3
<i>Lactuca serriola</i> + <i>Mj</i>	5.87 a	185.0	5.00 a	278.8
<i>Pulicaria crispa</i> + <i>Mj</i>	4.84 bc	135.0	3.86 d	192.4
<i>Reseda muricata</i> + <i>Mj</i>	4.55 bc	120.9	3.09 ef	134.1
<i>Salsola imbricata</i> + <i>Mj</i>	3.71 de	80.1	2.73 fg	106.8
<i>Trichodesma africanum</i> + <i>Mj</i>	4.47 cd	117.0	3.67 de	178.0
<i>Zygophyllum migahidii</i> + <i>Mj</i>	3.13 ef	51.9	1.68 hi	27.3

- Data are averages of five replicates.

- Values within each column followed by the same alphabetical letter(s) are not significantly different according to Fisher's protected LSD at $P= 0.05$.

* Increase (%) = $(G_T - G_N / G_N) \times 100$, where G_T = growth of the treatment and G_N = growth of the nematode check.

These results are in agreement with those given by Ibrahim and Ibrahim (2000), who found that soil treatment with powders of the air-dried brown alga *Botryocladia cappilaceae*, leaves of *Psidium guajava*, and cattle and chicken manures at 2% (w/w) gave a relative efficacy in suppressing *M. incognita*-infection on common beans approximately the same as the nematicide Carbofuran 10% applied at 0.2g/pot.

It is worth mentioning that application of these plant powders on a large scale under field or greenhouse conditions may have two constraints. First, we need great amounts of these plants and this is, sometimes, unavailable. Second, some species of wild plants, especially those grown on sludge or waste materials, have an ability to accumulate certain heavy metals toxic to plants such as Cu, Zn, Pb, Cd, Cr and Ni (Porębska and Ostrowska, 1999). Therefore, the adverse effect of *Z. migahidii* on the growth of common beans in the current study probably attributed to the presence of some heavy metals in its tissues. Studies of Taia and

El-Ghanem (2004) confirmed the presence of significant concentrations of Cu and Zn among mineral contents of *Z. migahidii* wildly grown in Riyadh, Saudi Arabia. On the other hand, some wild plants have allelopathic and toxic effects on other plants. It was found that shoot aqueous extract of *A. hierochuntica* inhibited seed germination, seedling growth, and cell division of certain plant species under laboratory conditions (Hegazy *et al.*, 1990). This observation may explain the phytotoxicity symptoms of *A. hierochuntica* appeared on common beans in the present study.

Lastly, wild plants with effective nematicidal activity in the current study should be subjected to further research works to investigate chemical structure of their bioactive constituent(s) with nematicidal activity, their mode(s) of action, and possibility of formulation.

We believe that natural wild vegetations still have a lot of hidden secrets and great benefits to human phytotherapy and nematode management.

Table 3. Phytochemical constituents of the studied wild plants that responsible for their biological activities and may involve in their nematicidal activity

Wild plant / Family	Phytochemical constituents	Reference(s)
<i>Achillea fragrantissima</i> (Asteraceae)	Glaucolides, sesquiterpene lactones, flavonoids, alkaloids, tannins, saponins, phenolic compounds and essential oil.	Abel-Mogib <i>et al.</i> , (1989) Al-Gaby and Allam, (2000) Al-Nowaihi <i>et al.</i> , (2005)
<i>Anastatica hierochuntica</i> (Brassicaceae)	Tannins, sterols, terpenes, flavonoids, alkaloids, saponins, resins, phenols and glycosides.	Mohamed <i>et al.</i> , (2000) Mohamed <i>et al.</i> , (2009)
<i>Brassica sinaica</i> (Brassicaceae)	Not available in the literature, but generally <i>Brassica</i> species contain glucosinolates and isothiocyanates. In addition, phenols and ascorbic acid in their tissues may compliment the activity of glucosinolates.	Chitwood, (2000) Antonious <i>et al.</i> , (2009)
<i>Calotropis procera</i> (Asclepidiaceae)	Alkaloids, cardiac glycosides, tannins, saponins, flavonoids, sterols and/or triterpenes.	Mossa <i>et al.</i> , (1991) Tanira <i>et al.</i> , (1994)
<i>Cleome rupicola</i> (Capparaceae)	Saponins, flavonoids, tannins, terpenes and/or sterols.	Tanira <i>et al.</i> , (1994)
<i>Ducrosia anethifolia</i> (Apiaceae)	Prenylated furanocoumarin pangelin and essential oil consisted of 20 compounds with majority of α -pinene (12.4%), n-decanal (70.1) and dodecanal (5.4%).	Janseen, <i>et al.</i> , (1984) Stavri <i>et al.</i> , (2003) Hajhashemi <i>et al.</i> , (2010)
<i>Fagonia bruguieri</i> (Zygophyllaceae)	Diterpenes, flavonoids and phenolics.	Abdel-Kader <i>et al.</i> , (1993) Tawaha <i>et al.</i> , (2007)
<i>Lactuca serriola</i> (Asteraceae)	Alkaloids, flavonoids, saponins and sesquiterpene lactones.	Mojab <i>et al.</i> , (2003) Michalska <i>et al.</i> , (2009)
<i>Pulicaria crispata</i> (Asteraceae)	Sesquiterpene lactones, kaurane glycosides, sesquiterpenes, monoterpenes, flavonoids, tannins, sterols, saponins, essential oil and phenolics.	Abdel-Mogib, <i>et al.</i> , (1990) Ahmed <i>et al.</i> , (2010) Liu <i>et al.</i> , (2010)
<i>Reseda muricata</i> (Resedaceae)	Flavonoids and phenolic acids.	El-Sayed <i>et al.</i> , (2001)
<i>Salsola imbricata</i> (Chenopodiaceae)	Alkaloids, coumarins, saponins, sterols and/or terpenes.	Al-Saleh, <i>et al.</i> , (1997)
<i>Trichodesma africanum</i> (Boraginaceae)	Essential oil, steroids, coumarins, flavonoids, phenolics, alkaloids and glycosides.	Abd El-Moaty, (2009)
<i>Zygophyllum migahidii</i> (Zygophyllaceae)	Not available in the literature, but generally <i>Zygophyllum</i> species contain glycosides and triterpenoid saponins in their tissues.	Hassanean <i>et al.</i> , (1993) Elgamal <i>et al.</i> , (1995)

ACKNOWLEDGEMENT

The authors highly appreciate cooperation and efforts of Dr. Jacob Thomas, Herbarium of Dept. of Botany and Microbiology, College of Science, King Saud University, Riyadh, for identifying species of the studied wild plants.

REFERENCES

Abd El-Moaty, H. I. (2009). Active constituents and antimicrobial activity of *Trichodesma africanum* (L.) R. Br. var. *heterotrichum* Bornm. & Kneuck. Bull. Fac. Agric. Cairo Univ. 60 (4): 357-365.

Abdel-Kader, M. S.; A. A. Omar; N. A. Abdel-Salam and F. R. Stermitz. (1993). Erythroxan diterpenes and flavonoids from *Fagonia bruguieri*. Phytochemistry 33 (3): 718-720.

Abdel-Mogib, M.; J. Jakupovic; A. M. Dawidar; M. A. Metwally and M. Abou-Elzahab. (1989). Glaucolides from *Achillea fragrantissima*. Phytochemistry 28 (12): 3528-3530.

Abdel-Mogib, M.; J. Jakupovic; A. M. Dawidar; M. A. Metwally and M. Abou-Elzahab. (1990). Sesquiterpene lactones and kaurane glycosides form *Francoeuria crispa*. Phytochemistry 29 (8): 2581-2584.

- Abdel-Sattar, E.; L. Maes and M. M. Salama. (2010). *In vitro* activities of plant extracts from Saudi Arabia against malaria, leishmaniasis, sleeping sickness and Chagas disease. *Phytotherapy Research* 24 (9): 1322-1328.
- Ahmed, E. M.; B. Y. M. Nour; Y. G. Mohammed and H. S. Khalid. (2010). Antiplasmodial activity of some medicinal plants used in Sudanese folk-medicine. *Environ. Health Insights* 4: 1-6.
- Al-Doghairi, M. A. and E. A. Elhag. (2003). Effect of *Rhazya stricta*, *Calotropis procera*, and *Francoeuria crispa* on larvae and eggs of *Culex pipiens*. *J. Herbs, Spices and Medicinal plants* 10 (2): 25-33.
- Al-Gaby, A. M. and R. F. Allam. (2000). Chemical analysis, antimicrobial activity, and the essential oils from some wild herbs in Egypt. *J. herbs, Spices and Medicinal Plants* 7 (1): 15-23.
- Al-Hazmi, A. S.; F. A. Al-Yahya and A. T. Abdul-Razig. 1995. Occurrence, distribution and plant associations of plant nematodes in Saudi Arabia. *Research Bulletin No. 52*, Department of Plant Protection, Agricultural Research Center, College of Agriculture, King Saud University.
- Al-Nowaihi, A. M.; A. M. A. Osman; T. A. M. Abdel-Raziek and T. M. El-Lamey. (2005). Effect of some ecological factors on the chemical compounds in some xerophytes for enhancing their use. *J. Environ. Sci.* 10 (2): 373-374.
- Al-Rajhi, D. H.; A. S. Al-Hazmi; H. I. Hussein; A. A. M. Ibrahim; F. A. Al-Yahya and S. Mostafa. (1997). Nematicidal properties of *Rhazya stricta* and *Juniperus polycarpos* on *Meloidogyne javanica* in Saudi Arabia. *Alex. Sci. Exch.* 18 (2): 135-142.
- Al-Saleh, F. S.; A. Y. Gamal El-Din; J. A. Abbas and N. A. Saeed. (1997). Phytochemical and biological studies of medicinal plants in Bahrain: the family Chenopodiaceae- Part 2. *Inter. J. Pharmacognosy* 35 (1): 38-42.
- Al-Yahya, F. A.; A. A. M. Dawabah; A. A. El-Sherbiny and A. S. Kamel. (2005). Effect of some plant extracts on the root-knot nematode (*Meloidogyne javanica*). Abstracts of 2nd Symposium of Saudi Society of Agricultural Sciences, 22-24, February 2005.
- Antonious, G. F.; M. Bomford and P. Vincelli. (2009). Screening *Brassica* species for glucosinolates content. *J. Environ. Sci. Health, Part B*, 44 (3): 311-316.
- Bogdadi, H. A. A.; L. Kokoska; J. Havlik; P. Kloucek; V. Rada and K. Vorisek. (2007). *In vitro* antimicrobial activity of some Libyan medicinal plant extracts. *Pharm. Biol.* 45 (5): 386-391.
- Chaudhary, S. A. (1999). Flora of the Kingdom of Saudi Arabia, vol. 1 (Illustrated). Ministry of Agriculture and Water, Riyadh, 640 pp.
- Chaudhary, S. A. and A. A. Al-Jowaid. (1999). Vegetation of the Kingdom of Saudi Arabia. Ministry of Agriculture and Water, Riyadh, 689 pp.
- Chitwood, D. J. (2002). Phytochemical based strategies for nematode control. *Ann. Rev. Phytopathol.* 40: 221-249.
- Dahiya, R.S. (2003). Effect of few organic amendments on cereal cyst nematodes, *Heterodera avenae* in wheat. In proceedings of national symposium on biodiversity and management of nematodes in cropping systems for sustainable agriculture, Jaipur, India, 11 -13 November, 2003.
- Elgamal, M. H. A.; K. H. Shaker; K. Pöllmann and K. Seifert. (1995). Triterpenoid saponins from *Zygophyllum* species. *Phytochemistry* 40 (4): 1233-1236.
- El-Sayed, N. H.; N. M. Omara; A. K. Yousef; A. M. Farag and T. J. Mabry. (2001). Kaempferol triosides from *Reseda muricata*. *Phytochemistry* 57 (4): 575-578.
- Hajhashemi, V; M. Rabbani; A. Ghanadi and E. Davari. (2010). Evaluation of antianxiety and sedative effects of essential oil of *Ducrosia anethifolia* in mice. *Clinics* 65 (10):1037-1042.
- Hassanean, H. H.; E. K. Desoky and M. M. A. El-Hamouly. (1993). Quinovic acid glycosides from *Zygophyllum album*. *Phytochemistry* 33 (3): 663-666.
- Hegazy, A. K.; K. S. Mansour and N. F. Abdel-Hady. (1990). Allelopathic and autotoxic effects of *Anastatica hierochuntica* L. *J. Chem. Ecol.* 16 (7): 2188-2193.
- Holbrook, C. C.; D. A. Knauf and D. W. Dikson (1983). A technique for screening peanut for resistance to *Meloidogyne arenaria*. *Plant Dis.*, 57: 957-958.
- Hussey R. S. and K. R. Barker. (1973). A comparison of methods of collecting inocula of *Meloidogyne* spp., including a new technique. *Plant Dis. Reporter* 57, 1025-1028.
- Ibrahim, A. A. M. and I. K. A. Ibrahim. (2000). Evaluation of non-chemical treatments in the control of *Meloidogyne incognita* on common bean. *Pakistan J. Nematol.* 18 (1&2): 51-57.
- Ibrahim, I. K. A.; A. A. M. Ibrahim and M. A. Rezk. (1994). Biological and chemical control of root-knot nematodes on clover, corn and wheat. *Alex. J. Agric. Res.* 39 (3): 453-462.
- Janssen, A. M.; J. J. C. Scheffer; A. B. Svendsen and Y. Aynehchi. (1984). The essential oil of *Ducrosia anethifolia* (DC.) Boiss: Chemical composition and antimicrobial activity. *Pharmaceutisch Weekblad Scientific Edition* 6 (4): 157-160.
- Liu, L. L.; J. L. Yang and Y. P. Shi. (2010). Phytochemicals and biological activities of *Pulicaria* species. *Chemistry and Biodiversity* 7 (2): 327-349.
- Maghraby, A. S.; N. Shalaby; H. I. Abd-Alla; S. A. Ahmed; H. M. Khaled and M. M. Bahgat. (2010). Immunostimulatory effects of extracts of *Pulicaria crispa* before and after *Schistosoma mansoni* infection. *Acta Poloniae Pharmaceutica – Drug Research* 67 (1): 75-79.
- Michalska, K.; A. Stojakowska; J. Malarz; I. Dolezalova; A. Lebeda and W. Kisiel. (2009). Systematic implications of sesquiterpene lactones in *Lactuca* species. *Biochemical Systematics and Ecology* 37 (3): 174-179.

- Mohamed, A. A.; A. A. Khalil and H. E. S. El-Beltagi. (2009). Chemical compositions and antioxidant/antimicrobial activities of kaff maryam (*Anastatica hierochuntica*) and doum palm (*Hyphaene thebaica*) cultivated in Egypt. *Biyoloji Bilimleri Araştırma Dergisi* 2 (2): 71-79.
- Mohamed, M. K.; S. N. Guergues and E. A. Abdel-Rahim. (2000). Studies on the phytochemistry and antimicrobial activity of four plant species from Egypt. *Egypt. J. Microbiol.* 35 (2): 257-271.
- Mojab, F.; M. Kamalinejad; N. Ghaderi and H. R. Vahidipour. (2003). Phytochemical screening of some species of Iranian plants. *Iranian J. Pharma. Res.* 2 (2): 77-82.
- Mossa, J. S.; M. Tariq; A. Mohsin; A. M. Ageel; M. A. Al-Yahya; M. S. Al-Said and S. Rafatullah. (1991). Pharmacological studies on the aerial parts of *Calotropis procera*. *American Journal of Chinese Medicine* 19 (3-4): 223-231.
- Nandal, S. N. and D. S. Bhatti. (1990). Efficacy, persistence and field application potential of some weeds/shrubs for controlling *Meloidogyne javanica* on brinjal. *Nematol. Medit.* 18: 113-115.
- Oka, Y. (2010). Mechanisms of nematode suppression by organic soil amendments - a review. *Applied Soil Ecology* 44 (2): 101-115.
- Oka, Y.; B. H. Ben-Daniel and Y. Cohen. (2001). Nematicidal activity of powder and extracts of *Inula viscosa*. *Nematology* 3 (8): 735-742.
- Oka, Y.; S. Nacar; E. Putievsky; U. Ravid; Z. Yaniv and Y. Spiegel. (2000). Nematicidal activity of essential oils and their components against the root-knot nematode. *Phytopathology* 90 (7): 710-715.
- Oostenbrink, M. (1966). Major characteristics of the relation between nematodes and plants. *Mededelingen Landbouwhogeschool Wageningen* 66:1-46.
- Pirbalouti, A.G. (2009). Medicinal plants used in Chaharmahal and Bakhtyari districts of Iran. *Herba Polonica* 55 (2): 69-77.
- Ploeg, A. T. and J. J. Stapleton. (2001). Glasshouse studies on the effects of time, temperature and amendment of soil with broccoli plant residues on the infestation of melon plants by *Meloidogyne incognita* and *M. javanica*. *Nematology* 3 (8):855-861.
- Porębska, G. and A. Ostrowska. (1999). Heavy metal accumulation in wild plants: Implications for phytoremediation. *Polish J. Environ. Studies* 8 (6): 433-442.
- Quarles, W. (1992). Botanical pesticides from *Chenopodium*? *The IPM Practitioner* 14 (2): 1-11.
- Rahman, M. A.; J. S. Mossa; M. S. Al-Said and M. A. Al-Yahya. (2004). Medicinal plant diversity in the flora of Saudi Arabia 1: a report on seven plant families. *Fitoterapia* 75 (2): 149-161.
- Rao, M. S.; P. P. Reddy and S. M. Das. (1996). Effect of integration of *Calotropis procera* leaf and *Glomus fasciculatum* on the management of *Meloidogyne incognita* infesting tomato. *Nematol. Medit.* 24: 59-61.
- Salamah, A. A.; H. M. Hassan and T. M. Nassar. (1989). Antibacterial wild flowering plants in Saudi Arabia. *J. King Saud Univ., Vol.1, Science (No.1,2)*: 5-19.
- SAS. (1997). SAS/STAT User's Guide: Statistics, Version 6.12. SAS Institute Inc. Cary, NC., USA.
- Sathiyamoorthy, P.; H. L. Evgi; P. Schlesinger; I. Kedar; J. Gopas; Y. Pollack and A. G. Goldhirsh. (1999). Screening for cytotoxic and antimalarial activities in desert plants of the Negev and Bedouin market plant products. *Pharm. Biol* 37 (3): 188-195.
- Stavri, M.; K. T. Mathew; F. Bucar and S. Gibbons. (2003). Pangelin, an antimycobacterial coumarin from *Ducrosia anethifolia*. *Planta Med.* 69 (10): 956-959.
- Taia, W. K. and W. M. El-Ghanem. (2004). Effect of habitats on both the phenotypic characters and mineral contents of five wild species in El-Riyadh city. *Pak. J. Biol. Sci.* 7 (8): 1399-1403.
- Tanira, M. O. M.; A. K. Bashir; R. Dib; C. S. Goodwin; I. A. Wasfi and N. R. Banna. (1994). Antimicrobial and phytochemical screening of medicinal plants of the United Arab Emirates. *J. Ethnopharmacol.* 41 (3): 201-205.
- Tawaha, K.; F. Q. Alali; M. Gharabeih; M. Mohammad and T. El-Elimat. (2007). Antioxidant activity and total phenolic content of selected Jordanian plant species. *Food Chemistry* 104 (4): 1372-1378.

الملخص العربي

فعالية ثلاثة عشر نوعاً من النباتات البرية كمحسنات للتربة في مكافحة نيماتودا تعقد الجذور

على نباتات الفاصوليا بالململكة العربية السعودية *Meloidogyne javanica*

عمره علي الشربيني، فهد بن عبد الله اليحيى

سجلت مساحيق نباتات *T. imbricata* ، *C. rupicola* ، *S. imbricata* ، *Z. migahidii* ، *africanum* أعلى نسبة خفض في المقاييس المرضية المذكورة تراوحت بين 82.0 - 96.4٪، بينما كانت أقل نسبة خفض (40.4 - 33.9٪) لمسحوق نبات *F. bruguieri* . وقد تسببت المعاملة بمبييد كاربوفيلوران 10٪ في خفض كبير في مقاييس الإصابة تراوح بين 88.0-90.8٪ . من ناحية أخرى أدت معظم المعاملات إلى زيادة معنوية في نحو نباتات الفاصوليا تراوحت بين 106.8-188.3٪ في المجموع الخضري ، وكذلك بالنسبة للمجموع الجنسي مقارنة بالنباتات المصابة بالنيماتودا، وكذلك بالنسبة للنباتات المعاملة بمبييد النيماتودي والأخرى السليمة. وعلى النقيض من ذلك، لوحظت أعراض سمية نباتية قليلة على نباتات المعاملات بمساحيق كلاً من النباتين *Z. migahidii* ، *A. hierochuntica* .

حققت مساحيق نباتات *T. imbricata* ، *C. rupicola* ، *S. imbricata* ، *Z. migahidii* ، *africanum* فعالية إبادية للنيماتودا تراوحت نسبتها بين 66.5 - 269.1٪ من فعالية مبييد كاربوفيلوران 10٪.

تعتبر تلك الدراسة بمنطقة التقرير الأول عن نباتات برية جديدة بالململكة العربية السعودية لها تأثير إبادي لنيماتودا تعقد الجذور، والتي ربما تكون مصدراً لمبيدات نيماتودية بديلة آمنة على البيئة والإنسان.

تمت دراسة فعالية إضافة مساحيق المجاميع الحضرية لثلاثة عشر نوعاً من النباتات البرية بالململكة العربية السعودية، كمحسنات للتربة، لمكافحة نيماتودا تعقد الجذور *Meloidogyne javanica* على نباتات الفاصوليا (صنف Strike) تحت ظروف الصوبة الزجاجية (درجة حرارتها $35\pm5^{\circ}\text{M}$) . النباتات المختبرة هي كالتالي: *Achillea* ، *Brassica* ، *Anastatica hierochuntica* ، *fragrantissima* ، *Cleome rupicola* ، *Calotropis procera* ، *sinaica* *Lactuca* ، *Fagonia bruguieri* ، *Ducrosia anethifolia* *Salsola* ، *Reseda muricata* ، *Pulicaria crispa* ، *serriola* *Zygophyllum* ، *Trichodesma africanum* ، *imbricata* *migahidii* . تم خلط المساحيق النباتية بالتربة الملوثة بالنيماتودا بتركيز 0.3٪ (وزن/وزن) ، قبل الزراعة بإسبوع . كانت هناك معاملة- للمقارنة- بمبييد النيماتودي كاربوفيلوران 10٪ (0.1 جم/الأصيص)، إضافة إلى معاملتين آخرتين كشواهد أحداها لنباتات سلية والأخرى لنباتات مصابة بالنيماتودا فقط. أظهرت النتائج أن جميع المساحيق النباتية المستخدمة أدت إلى خفض معنوي في أعداد كل من العقد الجندرية، كتل البيض، عدد البيض النهائي، وقيم عامل تكاثر النيماتودا المختبرة على جذور نباتات الفاصوليا، مقارنة بالنباتات المصابة بالنيماتودا فقط.