

Nematicidal Potentialities of Alginate Controlled-Release Formulations of Aldicarb and Cadusafos on The Root-Knot Nematode *Meloidogyne incognita* Infecting Tomato

Hamdy R. Soltan,¹ A. S. Marei and A.Mousa²

ABSTRACT

Two separate pot experiments was conducted to evaluate the nematicidal activity of the alginate controlled release (CR) formulations of aldicarb (FA₃, FA₆ and FA₉) and cadusafos (FC₃, FC₆ and FC₉) versus the corresponding granule formulations against the root-knot nematode *Meloidogyne incognita* infecting tomato under green house conditions. The results indicated that all the tested formulations significantly reduced the number of galls, egg masses, and 2nd juveniles in the soil as compared with the untreated pots (control). On the other hand, the CR formulations of aldicarb and cadusafos could be reduced the galls, egg masses, and 2nd juveniles in soil significantly compared with their granular formulations. The percentage reduction in nematode galls for all the CR formulations of aldicarb and cadusafos were ranged from (95.1% to 96.04%) and (87.1% to 88.03%) for aldicarb and cadusafos, respectively. Whereas, the lowest percentage of reduction of the number of tomato galls was recorded with aldicarb and cadusafos granule formulation (61.3% and 66.71%), respectively. Also, the maximum reduction of the number of 2nd juveniles in the soil was achieved by the FA₃ (86.73%) and FC₃ (89.08%), while for its corresponding granule was (59.05% and 46.23%). It could be noted that the nematicidal efficacy of the CR formulations against *M. incognita* reached to approximately 1.77 folds by FA₃ and 1.93 folds by FC₃ comparable to the commercial formulations. The possible effects of different aldicarb formulations residues on the chemical constituents of tomato fruits were determined and recorded at the end of the experiment (72 days). The results obtained indicating that all the aldicarb and cadusafos formulations had significantly decreased of the β-carotene, lycopene, and ascorbic acid content of tomato fruits comparable to the untreated pots. No significant differences were found between all the aldicarb formulations and the control on the percentage of the total soluble solid (T.S.S.) whereas there were significantly increased in the percentage of T.S.S. of tomato plant treated with the CR cadusafos formulations. The FA₃ and FC₃ could be applied on tomato plant to control nematode with a minimum side effect on the quality control parameters such as lycopene, T.S.S. and ascorbic acid content of tomato juice compared with the commercial formulations.

Key word: controlled release formulations, root-knot nematode, plant chemical constituent.

INTRODUCTION

Root-knot nematodes (*Meloidogyne* spp.) are among the most economically damaging genera of plant parasitic nematodes on horticultural and field crops, causing an estimated US\$100 billion loss globally on an annual basis (Oka *et al.*, 2000). *Meloidogyne incognita* is one of the most harmful root-knot nematode species, which infects a wide range of vegetable crops in Egypt (Ibrahim *et al.*, 2000). Root-knot nematodes infection of tomatoes limits fruit production by 30% (Anwar and McKenry, 2010). Among various control measures, use of chemicals has been considered as an effective strategy for control of plant parasitic nematodes when other methods like cultural practices, resistant varieties and bio-control agents are unable to protect crops from these pests (Randhawa *et al.*, 2001; Sakhuja and Jain, 2001). Currently various fumigants and non-fumigants are available in the market against these pests (Rich *et al.*, 2004). Granular non-fumigant nematicides are more easily applied and safer for farmers compared with fumigants (Lamberti *et al.*, 2000). Among the most extensively used non-fumigant granular are aldicarb and cadusafos which are carbamate and organophosphate based nematicides. Although chemical nematicides are effective, easy to apply and show rapid effects, they have begun withdrawn from the market owing to concerns about public health and environmental safety (Rich *et al.*, 2004). Controlled-release formulations technology offers the potential to reduce the environmental losses of pesticides and hence may be increased their efficacy, reduce their side effect to non-target organisms and reduce their harmful effect on growth and nutrient contents of crops. Biodegradable and synthetic matrices for carbofuran formulations were assessed against *M. incognita* infecting tomato by Radwan *et al.* (2006). Alginate gels are biodegradable and pesticides can be easily incorporated into the

¹Department of Applied Entomology, Faculty of Agriculture, Alexandria University, Egypt

²Agriculture Research Centre Sabahia Research Station, Alexandria Central Pesticide Research Lab

Received February 4, 2015, Accepted March 8, 2015

matrix using an aqueous system at ambient temperature (Soltan, 1991; Soltan 1996 and Soltan *et.al.*, 2014). For all the reasons mentioned above, alginate gels were selected as matrices for preparation of new controlled-release formulations of aldicarb and cadusafos nematicides. The objective of this research was to investigate if by using aldicarb or/and cadusafos in the form alginate controlled-release formulations, its efficacy vis-à-vis *M. incognita* control aspscoated tomato plants could be improved by evaluating the nematicidal efficacy expressed as the percent reduction of total number of galls, egg masses and 2nd juveniles nematode in the soil. The chemical constituents of tomato fruits were determined that may be affected by the aldicarb and cadusafos released from the different formulations

MATERIALS AND METHODS

Preparation on controlled – release formulations of aldicarb and cadusafos:-

Different types of alginate controlled-release formulations (CRF.s) for each nematicide (aldicarb and/or cadusafos) were prepared in the present study according to the method described by Soltan *et.al.* 2014. Among the CRF's prepared, FA₃, FA₆ and FA₉, for aldicarb and FC₃, FC₆ and FC₉, for cadusafos were tested. Commercial formulation (15G) for aldicarb and 10G for cadusafose was provided by Agrochem Company, Egypt, and FMC Co., U.S.A, respectively.

Nematicidal activity of aldicarb and cadusafos formulations against *Meloidogyne incognita* on tomato seedlings

Plastic pots (20 cm in diameter and 25 cm deep) were filled with 3Kg sand soil. Two tomato seedling (*Lycopersicon esculentum* L. var. strain B), three weeks old, were transplanted. Recommended practices for fertilization and irrigation were followed and the pots were left for one week before treatments in order to acclimate the tomato seedling under the greenhouse condition. After planting, treatments were arranged in a randomized complete block design with 6 replications for each treatment in addition to the control. Pots were all inoculated with the suspension of 5000 2nd juveniles of *M. incognita* per pot. Aldicarb formulations (FA₃, FA₆, FA₉ and 15G) and cadusafos formulations (FC₃, FC₆, FC₉, and 10G), 3 days latter were applied on the surface layer of the soil at the recommended rate (1.5mg and 2.4mg a.i. /kg soil) for aldicarb and cadusafos, respectively. The efficiency of the formulations tested was judged, seven weeks after nematicide formulations treatment, by determining the number of galls per root system, egg masses and 2nd juveniles' nematode in the soil. The number of gall per root system of plant was

accounted visually. The number of egg masses per root system was counted after staining with phloxin B (prepared by dissolving 0.12 g phloxin B per liter of water). The galled roots were placed in this solution for 15-20 minutes. The roots were gently rinsed in tap water. The egg masses were stained red and counted directly. On the other hand, the 2nd juveniles per 250 g soil were extracted according to Baermann plate technique (Goodey, 1963). The soil samples were suspended by tap water at least three times, and then screened through 200, 325 mesh sieve, respectively. The juveniles were washed and decanted from 325 mesh sieve to dish equipped with a tissue paper, and were left for 24 hours before counting under microscope.

Impact of aldicarb and cadusafos formulations on some internal quality parameters of tomato fruits:

Internal quality parameters of treated and untreated tomato fruits were measured at the end of experiment such as β -carotene, lycopene, ascorbic acid, and total soluble solid (T.S.S.). The method described by AOAC (1965) was followed to extract β -carotene and lycopene. The β Carotene color was measured at 450 nm against blank according to the method described by Umiel and Gabelman (1971) and lycopene measured at 503 nm according to the method described by Rangama(1977). Ascorbic acid was measured according to the method of AOAC (1984), Extraction and determination of total soluble solid (T.S.S.) in tomato fruits according to the method reported by Pearson (1979).

Statistical analysis:

Data were analyzed by analysis of variance and mean comparison by least significant difference (L.S.D.) measured at the probability level of 0.05 using SAS statistical software.

RESULTS AND DISCUSSION

Nematicidal activity of aldicarb and cadusafos formulations against *Meloidogyne incognita* on tomato plants:

The data in Tables (1 and 2) indicated that all the aldicarb formulations significantly reduced the number of root galling and the larval number of *M. incognita* as compared with the untreated pots. No significant differences among alginate controlled release formulations were found in the number of galls which were ranged from 87.1% to 95.1%. The granular formulation (15G) was the lowest effective formulation (61.3%) in this concern. The numbers of galls /root system recorded were 11.17, 18.33, and 29.5 for FA₃, FA₆, and FA₉, respectively. Whereas, for aldicarb 15G was 88.1 and control was 227.83.

Table 1. Effect of different aldicarb formulations on *Meloidogyne incognita* infecting tomato under greenhouse conditions

Treatment	*Mean number of galls/root system	Mean number of egg masses /root system	Mean number of 2 nd juveniles /250 g soil
Control	227.8±37 c	32400±4.51 d	668.17±1.37 d
FA ₃	11.3±7.8 a	7185±4.0 a	86.6±0.8 a
FA ₆	18.3±6.64 a	10410.7±4.51 ab	126.53±0.94 a
FA ₉	29.5±4.98 a	12663.7±6.67 b	207.3±0.94 b
15G	88.1±23.04 b	17136.7±3.8 c	273.61±1.0 c

*Each value is a mean ±SD of six replicates

Values followed by the same letter are not significantly different at $p \geq 0.05$.

Table 2. % Reduction in nematode parameters as affected by different aldicarb formulations

Treatment	Effectiveness on galls (%)	Effectiveness on egg masses (%)	Effectiveness on 2 nd juveniles/ 250g soil (%)
Control	-	-	-
FA ₃	95.1	77.82	86.73
FA ₆	91.95	67.87	80.76
FA ₉	87.1	60.95	68.87
15G	61.3	47.11	59.05

It should be cited that the CR formulations may be extended the nematicidal efficacy of aldicarb on tomato plant than its granular formulation. It can be concluded that there were negative correlation between the release rate of aldicarb formulations and the mean total number of galls/root of tomato seedlings, which in turn lead to increasing in tomato yield. In addition, the CRFs highly reduced the mean total number of egg masses comparable to the granular formulation. The number of egg masses reduction expressed as percentage of the total number of egg masses were ranged from 60.95% to 77.82% for CRFs and 47.11% for the granular formulation. Also, the controlled release formulations were the most effective formulations in the percent reduction on juveniles into galls and were ranged from 61.7% to 83.96%. Whereas the granular formulation considered the lowest effective one in this respects (44.9%). Deleon, (1984) reported that aldicarb decreased the population of *M. incognita*, by more than 57%. There was no significant difference on the mean of total egg asses into galls between FA₃ and FA₆ as well as between FA₆ and FA₉. Concerning to the number of 2nd juveniles 250g soil, the data showed that FA₃ and FA₆ were the most effective in reduction of the number of 2nd juveniles. Generally, it can be concluded that the highest of nematicidal efficacy was caused by the controlled-released formulations FA₃ (86.73%), FA₆ (80.76%), FA₉ (68.87%) whereas, the granular formulation gave the lowest effect (59.05%) according to the total number of 2nd juveniles / 250 gm soil. These results are in agreement with those obtained by El-Shoura *et al.*, (1992). They found that the controlled-

release formulations of carbofuran were the most effective treatment in reducing the *M. javanica* infecting tomato plant. Also, Parlaza *et al.*, (1979) found that aldicarb caused a marked reduction of *Meloidogyne spp.* population on carrot.

Data presented in Tables (3 and 4) indicated that cadusafos controlled-release formulations significantly reduced the number of root galls and 2nd juveniles in the soil than that of its commercial granule formulation. Radwan *et al.* (2012) reported that cadusafos granule was the least effective compound tested and caused 77.51 and 86.63% reduction in root galling and J₂ population in the soil, respectively.

Among the CR formulations, no significant deference in the number of galls and 2nd juveniles in the soil between FC₃ and FC₆. The percent reductions in nematode galls for all the controlled-release formulations were ranged from 88.03 to 96.04%. On the other hand, the lowest percent reduction in mean total number of galls was recorded associated tomato seedlings treated with the commercial formulation (66.71%) comparable to untreated treatment. Ibrahim (1994) found that the cadusafos applied at the rate (15-45 mg/pot) was totally suppressed the root-knot nematode on tomato plant.

Once again, according to the number of egg masses the result cited that the formulations tested were suppressed the mean total number comparable to the control and with a wide range of efficacy based on the formulation types.

Table 3. Effect of different cadusafos formulations on *Meloidogyne incognita* infecting tomato under greenhouse conditions

Treatment	*Mean number of galls/root system	Mean number of egg masses /root system	Mean number of 2nd juveniles /250 g soil \pm S.D.
Control	227.8 \pm 37 d	32400 \pm 4.51 d	668.17 \pm 1.37 d
FC ₃	9 \pm 4.81 a	6626.7 \pm 3.51 a	72.94 \pm 0.7 a
FC ₆	16.2 \pm 4.16 a	9800 \pm 4.58 ab	104.5 \pm 0.77 a
FC ₉	27.2 \pm 4.9 b	11176.7 \pm 3.51 b	184.25 \pm 1.34 b
10G	75.8 \pm 18.5 c	15263.3 \pm 2.1 c	239 \pm 0.92 c

*Each value is a mean \pm SD of six replicates

Values followed by the same letter are not significantly different at $p \geq 0.05$.

Table 4. % Reduction in nematode parameters as effected by different cadusafos formulations

Treatment	Effectiveness on galls (%)	Effectiveness on egg masses (%)	Effectiveness on 2nd juveniles/ 250g soil (%)
Control	-	-	-
FC ₃	96.04	79.55	89.08
FC ₆	92.86	69.75	84.36
FC ₉	88.03	65.5	72.42
10G	66.71	52.89	46.23

Generally, the CR formulations were more effective in reducing the mean total number of eggs than the granular formulation. In addition, the results indicated that the percent reduction in number of eggs into galls were ranged from 65.5% to 79.55% for CR formulations whereas, the percentage was 52.89% for commercial formulation.

The same trend was obtained concerning the mean total number of all juveniles into galls. The higher reduction was obtained from the roots treated with the CR formulations, mainly FC₃ (88.7%) and the lowest reduction was obtained from pots treated with the granule formulation (50.41%) compared to untreated pots. However, the efficacy of cadusafos formulations tested could be arranged descending as follow: FC₃-FC₆>FC₉>10G according to all parameters used to evaluate the efficacy of cadusafos formulations. These data was supported by those data obtained during the present course of study that reported the highest release rate, residue levels in soil as well as residue levels in the root zone of tomato plant was from the pots treated with FC₃ while, the lowest from the granular formulation. Also, it should be noted that the decreasing in the mean total number of eggs and the mean total number of all juveniles into gall, indicated that the alginate incorporated cadusafos might be improved its toxicity behavior from non-persistence nematicide to persistent one and/or improving the penetration of cadusafos into nematode galls.

Impact of aldicarb and cadusafos formulations on some internal quality parameters of tomato fruits:

Pesticide residues may cause no visible injury to the plants on which they are used. Some of the less apparent injurious effects of pesticides on plants are interference with biochemical and physiological processes in it. Any of this interference may temporary or permanently retard the growth of the plant and then crop production (yield), may lower its nutrition value, may even prevent its use as food, and may affect its quality characterization (Hance, 1981). The quality of plant and its validity for edible use by consumer depends on the contaminations with pesticide residues. This role of pesticides in altering the nutrient levels in the plants should be considered when using chemical available as potential pesticide. The effect of the aldicarb formulations tested on the quality parameters in tomato to juice was recorded and the data were presented in Table (5). The results indicated that all the aldicarb formulations had significantly decreased of the β -carotene, lycopene and ascorbic acid content comparable to the untreated pots in tomato fruits. No significant difference was found among all the aldicarb formulations and the control on the percentage of the total soluble solid (T.S.S.).

Also, no significant differences were found among the controlled-release formulations (CRFs) on the β -Carotene content, and ascorbic acid content whereas, there were significant difference between FA₉ and both FA₃ and FA₆ on the lycopene content.

Table 5. Effecte of Aldicarb Formulations Residue on some Quality Parameters of Tomato Fruits

Treatment	β -Carotene Content (ppm) ^c	Lycopene Content (ppm)	Ascorbic acid Content (ppm)	Total soluble solid (T.S.S.) (%) ^a
Control	57.64±15.2 _b	4.09±0.33 _b	44.8±3.5 _b	7.06±0.11 _a
FA ₃	36.36 ± 7.7 _b	3.46 ± 0.31 _b	31.5 ± 2.0 _b	7.13 ± 0.11 _a
FA ₆	30.96 ± 10.7 _b	3.55 ± 0.22 _c	33.83 ± 4.04 _b	7.13 ± 0.11 _a
FA ₉	41.24 ± 12 _a	3.78 ± 0.35 _a	33.25 ± 3.5 _a	7.2 ± 0.2 _a
15G	48.66 ± 18.8	3.32 ± 0.26	22.16 ± 2.0	7.15 ± 0.2

*Each value is a mean ±SD of six replicates.

Values followed by the same letter are not significantly different at $p \geq 0.05$.

On the other hand, the commercial formulation (15G) had the highest effect on the all quality control parameters, comparable to the CR formulations. These results are in disagreement with those reported by Baojaji and Mohajam (1978) who found that aldicarb had significant effect on each lycopene content and ascorbic acid content, but is in agreement with the results of the aldicarb in reducing β -Carotene content. From the above-mentioned results, it should be cited that the injury caused by the nematicide could be minimized by alteration the formulation type such as using the alginate controlled release formulation. Also among the alginate controlled release formulation, the FA₉ could be applied on tomato plant to control nematode with a minimum side effect on the quality control parameter such as the percentage of total soluble solids and ascorbic acid content of tomato juice. These parameters are very important in tomato storage period, tomato mature stage, nutrition value, palatability and the processing of different product from tomato juice.

These parameters are very important in tomato storage period, tomato mature stage, nutrition value, palatability and the process carried out of produce different products from tomato juice.

The effects of cadusafos released from the different formulations (CRFs and 10 G), which were uptaken from soil by tomato plant and translocated and reached to tomato fruits presented in Table (6). The statistical analysis of these data clearly indicated that there were

significant differences in the reducing of β -Carotene, lycopene, and ascorbic acid content between the tomato treated with cadusafos formulations and untreated one. In opposite, there were significant increases in the percentage of T.S.S. for the tomato plant treated with the CR cadusafos formulations as compared with the granular formulation as well as with the control. Among the CRFs, no significant differences were found in β -carotene content and total soluble solid. In addition, no significant difference was found in the ascorbic acid content between FC₃ and FC₆. Also, there was significant difference in the ascorbic acid content between FC₉ and the other tested formulations. On the other hand, the commercial formulation (10G) of cadusafos had the lowest value in total soluble solid and lycopene content comparable to the CR formulations. There was a highly significant difference between CF₉ and the other formulations in the ascorbic acid content. From all the results mentioned previously, we could conclude that the effect of cadusafos on the quality parameters tested were based upon the release rate of cadusafos which in turn based upon the formulation types. It is well established that certain organophosphorus compounds influence the chemical composition of the plant fruits to which they are applied. The insecticide, dimethoate did not influence T.S.S. and sugars of peppers. (Shahin *et al.*, 1989).

Table 6. Effecte of Cadusafos Formulations Residue on some Quality Parameters of Tomato Fruits

Treatment	β -Carotene Content (ppm) ^c	Lycopene Content (ppm)	Ascorbic acid Content (ppm)	Total soluble solid (T.S.S.) (%) ^a
Control	57.64±15.2 _b	4.09±0.33 _a	44.8±3.5 _a	7.06±0.11 _b
FC	26.95 ± 13.7 _b	3.47 ± 0.16 _b	33.25 ± 3.5 _a	7.33 ± 0.11 _b
FC ₃	38.7 ± 6.1 _b	3.66 ± 0.25 _b	35 ± 3.5 _b	7.36 ± 0.11 _b
FC ₆	37.65 ± 2.4 _a	3.64 ± 0.31 _a	39.08 ± 7.0 _a	7.35 ± 0.2 _a
10G	49.76 ± 5.7	3.35 ± 0.12	33.25 ± 3.5	7.1 ± 0.2

*Each value is a mean ±SD of six replicates.

Values followed by the same letter are not significantly different at $p \geq 0.05$.

Profenofos residues increased T.S.S. and acidity, but decreased the glucose, protein and ascorbic acid content of tomatoes (Ismail *et al.*, 1993). Radwan *et al.*, (1995) reported that pirimiphos-methyl residues appeared to have significant effect on the total soluble sugars and ascorbic acid content of tomato fruits and broad bean seeds. Consequently to obtain good results in pest chemical control in modern agriculture, the environmental factors, e.g. contamination of the treated edible crops with cadusafos residue and also the possible effect on the quality and nutritional values should be carefully considered to avoid problems which may arise as results of side effect of cadusafos.

REFERENCES

- Anwar, S.A. and M.V., McKrmry. 2010. Incidence and population density of plant-parasitic nematodes infecting vegetable crops and associated yield losses. *Pakistan J. Zool.*, 43: 327-333.
- AOAC. 1965. Official methods of analysis. 10th ed. pp.758.
- AOAC. 1984. Official methods of analysis of the Association of Official Analytical Chemists. Sidney Williams. 14thed. Association of Official Analytical Chemists, INC. Arlington, Virginia 22209 U.S.A.
- Baojaji, K.L., and R. Mohajam. 1978. Influence of some nematicides on the chemical composition of tomato fruits. *C.F. Horticultural Abst.* 48, 5:4677 a.
- Deleon, S.A.1984. Evaluation of Chemical products for the control of nematodes in coffee nurseries. *Revista Cafetalera.* 249:28-30.
- El-Shoura M.Y.; M.A, Radwan.; A.S.M. Marei and M. M. Abu Elamayem. 1992. Effect of aldicarb and carbofuran insect growth inhibitors and their mixture on root-knot nematicide on tomato plants. *J. Pest control & Environ. Sci.*, 4, (1):85-96.
- Goodey, J.B. 1963. Laboratory methods for work with plant and soil nematodes. *Minist. of Agric. Fisheries and Food: Tech. Bull.* 2. London: pp.44.
- Hance, R.J. 1981. Effects of pesticides on plant nutrition. *Residue Rev.*, 78:13-41.
- Ibrahim, A.A. M. 1994. Effect of cadusafos, *Paecilomyces lilacinus* and *Nemout ron* reproduction and damage potential of *Meloidogyn Javanica*. *Pakistan, J. of Nemato.*, 12, (2):141-147.
- Ibrahim, I.K.A.; Z.A. Handoo and A.A. El-Sherbiny. 2000. A survey of phytoparasitic nematodes on cultivated and non-cultivated plants in north western Egypt. *J. Nematol.*, 32: 478-485.
- Ismail, S.M.; H.M. Ali and R.A. Habiba. 1993. GC-ECD and GC-MS analysis of profenofos residues and biochemical effects in tomatoes and tomato products. *J. Agric. Food Chem.* 41: 610-615.
- Lamberti, F.; T.D. Addabbo., N. Sasanelli; Carella, A. and P. Greco. 2000. Chemical control of root-knot nematodes. *Acta Horticult.*, (ISHS), 532:183-188.
- Oka, Y.; H. Koltai; M. Bar-Eyal; M. E. I. Sharon; and Y. Spiegel. 2000. New strategies for the control of plant-parasitic nematodes. *Pest Management Science*, 56: 983-988.
- Pearson, D. 1979. The chemical analysis of foods. Seventh edition, Chapter Seven. pp. 233, Printed in Great Britain by T&A. Constable Ltd., Edinburgh.
- Perlaza, F.; R. Lopez and E. Vargas. 1979. Chemical control of *Meloidogyne spp.* and *Alternaria sp.* in Carrots (*Daucus Carota* L.). *Turrialba*, 29,(4):263-67. (*C.F.Chem. Abst.*, 1980, 92, 23:189, 192716 g).
- Radwan, M.A.; M.M. Youssef; A. Abd El-all; G.L.Henawy; and A.E. Marei. 1995. Residue levels of pirimiphos-methyl and chlorpyriphos-methyl on tomato and faba bean plants in relation to their impact on some internal quality parameters. *Alex. Sci. Exch.*,16 (3):389-404.
- Radwan, M.A.; S.A.A Farrag; M.M. Abu-El-Amayem; and N.S. Ahmed. 2012. Efficacy of some granular nematicides against root-knot nematode, *Meloidogyne incognita* associated with tomato. *Pakistan J. Nematol.*, 30:41-47.
- Radwan; M.A.; H.S. Ibrahim; S.A.A. S.M.I. Kassem and M.M. Abu-El-Amayem. 2006. Preliminary evaluation of the manggement of *Meloidogyne incognita* using biodegradable and synthetic matrices for carbofuran formulations. *Indian J. Nematol.*, 36 (1): 54-59.
- Randhawa, N.; P.K. Sakhujand and I. Singh. 2001. Management of root-knot nematode *Meloidogyne incognita* in tomato with organic amendments. *Plant Disease Research*, 16: 274- 276.
- Rangama, S. 1977. Manual of analysis of fruit and vegetable products. Tata McGraw-Hill Pub. Co. Ltd. New Delhi.
- Rich, J.R.; R. Dunn and J. Noling. 2004. Nematicides: Past and present uses. In: *Nematology: Advances and Perspectives*, Vol 2. Nematode Management and Utilization. (Eds.) Chen, Z.X.; S.Y. Chen and D.W. Dickson, CABI Publishing, Wallingford, UK.
- Sakhuja, P.K. and R.K. Jain. 2001. Nematode diseases of vegetable crops and their management. In: *Diseases of Fruits and Vegetables and their Management.* (Thind, T. S. ed.), Kalyani Pub., Ludhiana, India, pp: 474.
- Shahin, A.; E.M.R. Agwah and S.E. Mohamedin. 1989. Effect of dimethoate on greenhouse pepper and cucumber fruits quality and its residual behavior. The 7th Arab Pesticide conference, Tanta Univ., Sept. 11-12: 279-287.
- Soltan, H.R. 1991. Environmental studies on certain pesticides "Environmental Behaviour of the slow release formulations of Butachlor herbicides". PhD., Dept. Pesticides Chemistry, Fac. Of Agric., Alexandria University, Alex. Egypt.

- Soltan, H.R. 1996. Controlled release formulations of carbofuran based on alginate-kaolin-agriculture waste. *Alex. J. Agric. Res.*, 41 (1):241-252.
- Soltan, H.R.(2002). Colorimetric method for determination of the nematicides, cadusafos in aqueous medium at different temperature and PHvalues. *J.Adv. Agric. Res.*,7,1:183-92.
- Soltan, H.R.; A.S.M Marei and A. Mousa. 2014. Determination of the release rate profiles of aldicarb and cadusafos as alginate controlled- release formulations released in water by using newly modified colorimetric methods. *Alex. J. Agric. Res.* , 59 (2): 113-119.
- Umiel, N. and W.H. Gabelman. 1971. Analytical procedures for detecting carotenoids of carrot (*Daucus Carota* L.) roots and tomato (*Lycopersicum esculentum*) Fruits. *J. Amer. Soc. Hort. Sci.*, 96 (6):702-704

(,) FC₃ (,)

FA₃)

(FC₉ FC₆ FC₃)

(FA₉ FA₆

FA₃

FC₃

(, - ,)

FC₃ FA₃

(, - ,)

(, ,)

FA₃