

## **Comparative Toxicity of Four Formulations of Commercially Used Control Agents to The Red Flour Beetle, *Tribolium Castaneum* (Herbst) Under Laboratory Conditions**

Mohammed A. Al-Deghairi<sup>1</sup>

### **ABSTRACT**

Four formulations of commercially used control agents were tested under laboratory conditions on adults of the red flour beetle, *Tribolium castaneum*. The bioassays were carried out on wheat grains, at  $25 \pm 1$  °C and  $70 \pm 5\%$  RH. Compounds were applied at the dose rates of 250, 500, 750 and 1000 mg/L for Fenvalerate, 2500, 5000, 10000 and 20000 mg/L for Biofly, 25, 50, 100 and 200 mg/L for Neem and 25, 50, 75 and 100 mg/L for Diazinon. Adult mortality was assessed every day for 10 days and LC<sub>50</sub>, LC<sub>95</sub>, LT<sub>50</sub> and LT<sub>95</sub> were calculated. Mortality was 100% on wheat treated with 1000 mg/L of Fenvalerate, and 50, 75 and 100 mg/L of diazinon 10 days after exposure. Diazinon was the most effective against *T. castaneum* followed by Neem, Fenvalerate and then Biofly, where the recorded LC<sub>50</sub> values were 46.25, 78.30, 271.12 and  $3.05 \times 10^4$  mg/L after 4, 8, 7 and 6 days of exposure, respectively. Generally, for all tested compounds, the LT<sub>50</sub> values decreased as concentrations increased and Diazinon provided the shorter exposure time required for mortality. In conclusion, the biological method would not be as efficient as chemicals. However, the use of an effective plant extracts such as Neem on the stored grains would decrease the frequency with which insecticide was required. Thus, potentially remaining suitable pest control with a reduction in chemical residue. The present results encourage the use of Neem in control of stored product pests such as *T. castaneum* as alternative for synthetic insecticides or as a part in the integrated pest management programmes.

### **INTRODUCTION**

The red flour beetle, *Tribolium castaneum* (Herbst) is a serious pest of stored grain and grain products in many parts of the world. The beetles feed both as larvae and adults, and cause damage resulting in high economic losses. *Tribolium castaneum* secretes specific compounds called *p*-benzoquinones such as methyl-1,4-benzoquinone (MBQ) and ethyl-1,4-benzoquinone (EBQ) that are major components of defensive secretions used as repellents and irritants (Blum, 1981;

Howard, 1987; Eisner *et al.*, 1998), these benzoquinones are hazardous to human health.

The chemical control has focused on very few compounds as control agents for stored product insects. Currently there is renewed interest in developing reduced-risk low toxicity chemicals, including new formulations of insect growth regulators (IGRs), to replace older conventional products used in many agricultural systems, including post-harvest protection in areas where food is stored. Oberlander *et al.* (1997) reviewed many studies involving IGRs and stored-product insects. Most of them involved exposure of eggs, 1<sup>st</sup> instars, or adults on treated grains, in diet, or in glass vials, with subsequent measurements of progeny production. The efficacy of hydroprene (an IGR) on *T. castaneum*, was studied by Arthur (2003) and Arthur and Hoernemann (2004). One of the frequently used IGR is lufenuron ((*RS*)-1-[2,5-dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy) phenyl]3-(2,6-difluorobenzoyl) urea (Kedar *et al.*, 2008). This class of chemicals acts as chitin synthesis inhibitor (CSIs). Recently, CSIs have gained significant popularity due to their low mammalian toxicity and absence of mutagenic and teratogenic effects on warm-blooded animals. The residual analysis of lufenuron in wheat grains as well as the biological efficacy opens up a new vista for its possible use in protection of the stored food commodities (Kedar *et al.*, 2008).

Other chemical agents like synergized pyrethrins and the organophosphate dichlorvos are two insecticides that have historically been used for aerosol applications inside processing facilities and food warehouses. More recently, new pyrethroid and insect growth regulators have been registered as aerosols for use against stored-product insects, but most documented reports of efficacy describe tests done under laboratory conditions (Arthur and Gillenwater, 1990; Arthur, 1993). Recently methyl

<sup>1</sup>Plant Production & Protection Dept., College of Agric. & Vet. Medicine, P.O. Box 6622, Qassim University, Buraidah, Al-Qassim, Saudi Arabia  
Received July 13, 2008, Accepted August 6, 2008

bromide as an efficient fumigant to stored product insects was banned because it causes depletion of the ozone layer in the stratosphere.

Therefore, the control of red flour beetle has taken several approaches as mentioned by many scientists in that field. Su (1983) found that acetone extracts of *Piper nigrum* L. and the hexane extract of dry fruit of *P. guineese* was toxic to four species of stored product insects. Sighamony *et al.* (1984) also reported that acetone extracts of black pepper seeds were repellent to *T. castaneum* (Herbst). The essential oils of several spices like anise (*Pimpinella anisum* L.) and peppermint (*Mentha piperita* L.) were found to have fumigant toxicity to four major stored product pests, *Rhyzopertha dominica* (Fab.), *T. castaneum*, *Sitophilus oryza* (L.) and *Orzaephilus surinamensis* (L.) (Shaaya *et al.*, 1991).

The non-polar extracts of the flower buds of clove, *Syzygium aromaticum* (L.) and star anise (*Illicium verum* Hook f.) have insecticidal activity to *T. castaneum* and *S. zeamais* Motsch as well as suppress progeny production (Ho *et al.*, 1994 and 1995). Moreover, the essential oil of garlic was found to be potent against *T. castaneum* and *S. zeumais* (Ho *et al.*, 1996), while chopped garlic and garlic extracts were repellent to the two beetles species (Ho and Ma, 1995). Huang and Ho (1998), found that the essential oil of *Cinnamomum aromaticum*, was toxic to both *T. castaneum* and *S. zeamais* with contact fumigant and antifeedant activities.

The metabolite of the actinomycete *Saccharopolyspora spinosa* Mertz and Yao, Spinosad, is currently used on several stored products (Thompson *et al.*, 1997). It has low mammalian toxicity and it is very effective against a wide range of pest species (Cloyd and Sadof, 2000; Peck and McQuate, 2000). Recently, spinosad was registered for use in stored products in the USA (Subramanyam *et al.*, 2003), as an alternative to traditional grain protectants.

Due to the urgent need to develop safe, convenient, environmental and low-cost alternatives to control stored product pests, the present study was designed to assess the efficacy of four control agents which belong to different classes, namely Fenvalerate (pyrethroid), Diazinon (organophosphate), Neem (Natural product) and Biofly (Entomopathogenic fungus).

## MATERIALS AND METHODS

### Chemicals:

Fenvalerate, 20% EC (cyano(3-phenoxyphenyl) methyl 4-chloro- $\alpha$ -(1-methylethyl) benzene acetate; Neem (azadirachtin), 0.09% EC (dimethyl [2aR [2 $\alpha$ a,3 $\beta$ ,4 $\beta$ (1aR\*,2S\*,3aS\*,6aS\*,7S\*,7aS\*)],4a $\beta$ ,5 $\alpha$ ,7aS\*

,8 $\beta$ (E),10 $\beta$ ,10 $\alpha$ ,10b $\beta$ ]-10-(acetyloxy)octahydro-3,5-dihydroxy-4-methyl-8-[(2-methyl-1-oxo-2-butenyl)oxy]-4-(3a,6a,7,7a-tetrahydro-6a-hydroxy-7a-methyl-2,7-methanofuro[2,3-*b*] oxireno [*e*] oxepin-1a(2*H*)-yl)-1*H*,7*H*-naphtho[1,8-*bc*:4,4a-*c'*]difuran-5,10a(8*H*)-dicarboxylate; Diazinon, 60% EC (*O,O*-diethyl *O*-[6-methyl-2-(1-methylethyl)-4-pyrimidinyl] phosphorothioate) and Biofly, 100% (W/W) (Entomopathogenic fungus, *Beauveria bassiana*) were purchased from the local market.

### Insect rearing:

The red flour beetle *Tribolium castaneum*, used throughout the present work was maintained under laboratory conditions at 25  $\pm$  1°C and 70  $\pm$  5% relative humidity on wheat grains for several generations.

### Toxicity of control agents against *T. castaneum*

Tested concentrations were 250, 500, 750 and 1000 mg/L for Fenvalerate, 2500,5000,10000 and 20000 mg/L for Biofly, 25, 50, 100 and 200 mg/L for Neem and 25, 50, 75 and 100 mg/L for Diazinon. Concentrations were thoroughly incorporated in wheat grains using water as carrier solvent. The treated grains were kept for 24 hr. at 25°C till complete evaporation of the solvent before starting the experiments. Ten adults of *T. castaneum* were put on 25 gm of the treated wheat grains for 10 days or until 100% mortality was obtained. The data was subjected to probit analysis in order to determine the LC<sub>50</sub>, LC<sub>95</sub>, LT<sub>50</sub> and LT<sub>95</sub> values. All experiments were carried out in triplicates at 25  $\pm$  1°C and average 70  $\pm$  5% relative humidity.

### Statistical analysis

Data were calculated as mean  $\pm$  standard deviation (SD) analyzed using analysis of variance (ANOVA). Probability of 0.05 or less was considered significant. The statistical package of Costat Program (1986) was used for all chemometric calculations.

## RESULTS AND DISCUSSION

Four commercially used control agents, Fenvalerate, Neem, Biofly and Diazinon, were tested against the adult stage of *T. castaneum* (Tables1-4) under the laboratory conditions. The results indicated that the mean values of the percent mortality were significantly different. The average mortalities were 41.66, 47.99, 53 and 62% at 250, 500, 750 and 1000 mg/L of Fenvalerate, respectively (Table1). Less average mortality values were recorded in the case of Biofly with values being 10, 10, 21.70 and 23.33% at 2500, 5000, 10000 and 20000 mg/L, respectively (Table 2). In the case of Neem, the recorded average mortality values were 26.95, 34.33, 35.19 and 37.84 at 25, 50, 100 and 200 mg/L, respectively (Table 3). The highest average mortality values were recorded with Diazinon of 45.64,









57.61, 66.62 and 69.10% at 25, 50, 75 and 100 mg/L, respectively (Table 4). The 100% mortality was observed at 75 and 100 mg/L of Diazinon and 1000 mg/L of Fenvalerate after 6 and 10 days, respectively, compared with the control which showed less than 5% mortality in all cases. These results clearly indicate the high potential of Diazinon as a recommended treatment of wheat grains for controlling *T. castaneum* infestation.

As for the  $LT_{50}$  values for the adult stage of *T. castaneum*, they were found to be 6.50, 4.96, 4.04 and 3.15 days at 250, 500, 750 and 1000 mg/L of Fenvalerate (Table 5), while the  $LT_{50}$  values for Biofly were  $4.08 \times 10^3$  and 98.00 days at 10000 and 20000 mg/L, respectively (Table 6). The  $LT_{50}$  values for Neem were 11.50, 10.42, 9.21 and 7.75 days at 25, 50, 100 and 200 mg/L, respectively (Table 7). For Diazinon, it was found to have lower values of  $LT_{50}$  being 3.78, 2.74, 2.23 and 2.10 days at 25, 50, 75 and 100 mg/L, respectively (Table 8).

The  $LC_{50}$  values for the *T. castaneum* adult were found to be 46.25, 78.30, 271.12 and  $3.05 \times 10^4$  mg/L, while  $LC_{05}$  values were 304.64,  $2.25 \times 10^5$ ,  $4.20 \times 10^3$  and  $1.71 \times 10^{10}$  mg/L after 4, 8, 7 and 6 days of exposure for Diazinon, Neem, Fenvalerate and Biofly, respectively (Table 9). These results indicated that Diazinon was the most effective insecticide on *T. castaneum*, while the Biofly was the least.

### CONCLUSION

Storage of grains is part of the post-harvest practice through which food material passes on its way from fields to consumer. There is a continuous need to protect stored products against deterioration, especially loss of quality and quantity during storage. The use of pesticides is one means of preventing some losses during storage. However, the use of pesticides for storage pest control is very limited because of the strict requirement imposed on the use of synthetic insecticides on or near food (Padín *et al.*, 2002). The continuous use of chemical pesticides for control of stored-grain pests has resulted in serious problems such as resistance (Pacheco *et al.*, 1990). The present results demonstrated that Diazinon was most effective against *T. castaneum* followed by Neem, Fenvalerate and then Biofly. Furthermore, the efficacy of the insecticides against storage pest varies greatly after treatment (Suchita *et al.*, 1989 and Pinto *et al.*, 1997). Also, chemicals used for stored product pests, or as protectants, need also to be compared with the suitability and effectiveness of alternative control method. Non-chemical methods are attractive since they neither have chemical residues in the commodity nor do they cause resistance in insects. The public awareness and concern for environmental quality, has led to more focused attention on research

and development of biological agents, (Hidalgo *et al.*, 1998) and plant extract (Jbilou *et al.*, 2008) either as alternative or in integrated programs. A promising strategy with good potential to minimize the adverse effects of synthetic insecticides is the use of entomophagous fungi and other microbial control agents. The possibility of using fungal pathogens to control insects has been studied for many years but little attention has been paid to use of fungi as control agents against storage pests (Khan and Selman, 1988; Rodrigues and Partissoli, 1990; Adane, 1994; Adane *et al.*, 1996; Padín *et al.*, 1997; Hidalgo *et al.*, 1998; Moino *et al.*, 1998 and Padín *et al.*, 2002). The present results are in parallel with the results of Padín *et al.* (2002) who indicated that *B. bassiana* was not effective against *T. castaneum*, while *S. oryza* was very susceptible to *B. bassiana* isolate.

In general, the biological methods would not be as efficient as chemicals. However, the use of effective plant extracts such as Neem on the stored grain would reduce the need for synthetic insecticides. Thus, potentially remaining suitable pest control with a reduction in chemical residue. The present results encourage the use of Neem in control of stored product pests such as *T. castaneum* as alternative for synthetic insecticides or in the integrated programs.

### REFERENCES

- Adane, K. (1994). Microbial control of storage pests using the entomopathogenic fungus, *Beauveria bassiana* with special references to *Sitophilus zeamais* and *Callosobruchus chinensis*. M.Sc. Thesis, University of London, 93pp.
- Adane, K., Moore, D. and Archer, S.A. (1996). Preliminary studies on the use of *Beauveria bassiana* to control *Sitophilus zeamais* (Coleoptera: Curculionidae) in the laboratory. *Journal of Stored Products Research* 32, 105–113.
- Arthur, F. H. (1993). Evaluation of prallethrin aerosol to control stored product insect pests. *Journal of Stored Products Research* 29, 253–257.
- Arthur, F. H., (2003). Efficacy of a volatile formulation of hydroprene (Pointsource TM) to control *Tribolium castaneum* and *Tribolium confusum* (Coleoptera: Tenebrionidae). *Journal of Stored Products Research* 39, 205–212.
- Arthur, F. H. and Gillenwater, H.B. (1990). Evaluation of esfenvalerate aerosol for control of stored product insect pests. *Journal of Entomological Science* 25, 267.
- Arthur, F. H. and Hoernemann, C. K (2004). Impact of physical and biological factors on susceptibility of *Tribolium castaneum* and *Tribolium confusum* (Coleoptera: Tenebrionidae) to new formulations of hydroprene. *Journal of Stored Products Research* 40: 251–268.

- Blum, M.S. (1981). Chemical Defenses of Arthropods. Academic Press, New York, pp. 183–205.
- Cloyd, R.A. and Sadof, C.S. (2000). Effects of spinosad and acephate on western flower thrips inside and outside a greenhouse. *Hortechology* 10: 359–362.
- Costat Program (1986). Version 2, Cohort Software, Minneapolis, MN, USA.
- Eisner, T., Eisner, M., Attygalle, A.B., Deyrup, M. and Meinwald, J. (1998). Rendering the inedible edible: circumvention of a millipede's chemical defense by a predaceous beetle larva (Phengodidae). *Proc. Natl. Acad. Sci. USA* 95, 1108–1113.
- Hidalgo, E., Moore, D. and Le Patourel, G. (1998). The effect of different formulations of *Beauveria bassiana* on *Sitophilus zeamais* in stored maize. *Journal of Stored Products Research* 34, 171–179.
- Ho, S. H., Cheng, L. P. L., Sim, K. Y. and Tan, H. T. W. (1994). Potential of cloves (*Syzygium aromaticum* L.) Merr. and Perry as a grain protectant against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Postharvest Biology and Technology* 4, 179–183.
- Ho, S. H., Koh, L., Ma, Y., Huang, Y. and Sim, K. Y. (1996). The oil of garlic, *Allium sativum* L. (Amaryllidaceae), as a potential grain protectant against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Postharvest Biology and Technology* 9, 41–48.
- Ho, S. H. and Ma, Y. (1995). Repellence of some plant extracts to the stored product beetles, *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. Paper presented at the Symposium on Pest Management for Stored Food and Feed. SEMEO BIOTROP, Bogor, Indonesia, 5-7 September, 1995.
- Ho, S. H., Ma, Y., Goh, P. M. and Sim, K. Y. (1995). Star anise, *Illicium uerum* Hook f. as a potential grain protectant against *Tribolium castaneum* (Herbst) and *Sitophilus reamais* Motsch. *Postharvest Biology and Technology* 6, 341–347.
- Howard, R.W. (1987). Chemosystematic studies of the *Tribolium* (Coleoptera: Tenebrionidae): phylogenetic inferences from the defensive chemicals of eight *Tribolium* spp., *Palorus ratzeburgi* (Wissmann), and *Latheticus oryzae* Waterhouse. *Ann. Ent. Soc. Amer.* 80, 398–405.
- Huang Y. and S. H. Ho (1998). Toxicity and antifeedant activities of cinnamaldehyde against the grain storage insects, *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Journal of Stored Products Research* 34, 11–17.
- Jbilou, R.; Amri, H.; Bouayad, N., Ghailani, N.; Ennabili, A. and Sayah, F. (2008). Insecticidal effects of extracts of seven plant species on larval development,  $\alpha$  amylase activity and offspring production of *Tribolium castaneum* (Herbst) (Insecta: Coleoptera: Tenebrionidae). *Bioresource Technology* 99: 959–964.
- Kedar, C. A., Manjit, S.A., and Samindra, N. M. (2008). Development and application of a method for analysis of lufenuron in wheat flour by gas chromatography–mass spectrometry and confirmation of bio-efficacy against *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Chromatography B*, 861: 16–21
- Khan, A.R. and Selman, B.J. (1988). On the mortality of *Tribolium castaneum* adults treated sublethally as larvae with pirimiphos methyl, Nosema whitei and pirimiphos methyl- *N. whitei* doses. *Entomophaga* 33, 377–380.
- Moino A. Jr., Alves, S.B., and Pereira, R.M. (1998). Efficacy of *Beauveria bassiana* (Balsamo) Vuillemin isolates for control of stored-grain pests. *Journal of Applied Entomology* 122, 3
- Oberlander, H., Silhacek, D. L., Shaaya, E. and Ishaaya, I. (1997). Current status and future prospects of the use of insect growth regulators for the control of stored-product insects. *Journal of Stored Products Research* 33, 1–6.
- Pacheco, I.A., Sartori, M.R. and Taylor, R.W.D. (1990). Levantamento de resistência de insetos-pragas de gra armazenados a fosfina, no Estado de Sao Paulo. *Coletanea do Instituto de Tecnologia de Alimentos* 20, 144–154.
- Padín, S.B., Dal Bello, G.M., Vasicek, A.L. (1997). Pathogenicity of *Beauveria bassiana* for adults of *Tribolium castaneum* (Col.: Tenebrionidae) in stored grains. *Entomophaga* 42, 569–574.
- Padín, S.B., Padín, and Fabrizio, M (2002). Grain loss caused by *Tribolium castaneum*, *Sitophilus oryzae* and *Acanthoscelides obtectus* in stored durum wheat and beans treated with *Beauveria bassiana*. *Journal of Stored Products Research* 38, 69–74.
- Peck, S.L. and McQuate, G.T. (2000). Field tests of environmentally friendly malathion replacements to suppress wild Mediterranean fruit fly (Diptera: Tephritidae) populations. *Journal of Economic Entomology* 93, 280–289.
- Pinto A.R. Jr., Furiatti, R.S., Pereira, P.R.V.S. and Lazzari, F.A. (1997). Avaliac, ao de Inseticidas no Controle de *Sitophilus oryzae* L. (Coleoptera: Curculionidae) e *Rhyzopertha dominica* Fab. (Coleoptera: Bostrichidae) em Arroz Armazenado. *Anais da Sociedade Entomologica do Brasil* 26, 285–290.
- Rodrigues, C., and Partissoli, D. (1990). Patogenicidade de *Beauveria brongniartii* (Sacc.) Petch. e *Metarhizium anisopliae* (Metsch.) Sorok. e seu efeito sobre o gorgulho do milho e caruncho do feijao. *Anais da Sociedade Entomologica do Brasil* 19, 301–306.
- Shaaya, E., Ravid, U., Paster, N., Juven, B., Zisman, U. and Pissarev, V. (1991). Fumigant toxicity of essential oils against four major stored-product insects. *Journal of Chemical Ecology* 17, 499–504.
- Sighamony, S., Anees, I., Chandrakala, T. S. and Osmani, Z. (1984). Natural products as repellents for *Tribolium castaneum* Herbst. *International Pest Control* 26, 156–I 57.
- Su, H. C. F. (1983) Comparative toxicity of three Peppercorn extracts to four spices of stored product insects under laboratory conditions. *Journal of Georgia Entomological Society* 19, 190–199.

- Subramanyam, Bh., Toews, M.D. and Fang, L. (2003). Spinosad: an effective replacement for organophosphate grain protectants. In: Credland, P.F., Armitage, D.M., Bell, C.H., Cogan, P.M., Highley, E. (Eds.), Proceedings of the Eighth International Working Conference on Stored-Product Protection, 22–26 July 2002, York, UK. CAB International, Wallingford, Oxon, pp. 916–920.
- Suchita, M.G., Reddy, G.P.U., and Murthy, M.M.K. (1989). Relative efficacy of pyrethroids against rice weevil (*Sitophilus oryzae* L.) infesting stored wheat. Indian Journal of Plant Protection 17, 243–246.
- Thompson, G.D., Michel, K.H., Yao, R.C., Mynderse, J.S., Mosburg, C.T., Worden, T.V., Chio, E.H., Sparks, T.C. and Hutchins, S.H. (1997). The discovery of *Saccharopolyspora spinosa* and a new class of insect control products. Down to Earth 52, 1–5.

## الملخص العربي

# مقارنة التأثيرات السامة لأربعة من المركبات التجارية ضد خنفساء الدقيق الصدفية تحت الظروف المعملية

مُجد بن عبد العزيز الدغيري

أوضحت النتائج أن مبيد الدايازينون كان الأكثر فعالية ضد يرقات خنفساء الدقيق الصدفية يليه النيم ثم الفنفاليريت ثم البيوفلاي حيث بلغت قيم التركيز اللازم لقتل 50% بما يعادل 46.25 و 78.30 و 271.12 و  $10 \times 3.05$  مجم/لتر بعد 4 و 8 و 7 و 6 أيام من التعرض، على التوالي.

وبصفة عامة ولكل المركبات المختبرة فإن مقدار الزمن اللازم لقتل 50% من يرقات خنفساء الدقيق الصدفية تتناقص بزيادة التركيز و أن مركب الدايازينون قد احتاج لوقت أقل من التعرض لإحداث الموت.

ويمكن القول بأن استخدام المبيدات الميكروبية غير فعال مقارنة بالمبيدات الكيميائية، إلا أن استخدام النيم كمستخلص نباتي على الحبوب المخزونة قد يقلل من الاعتماد على استخدام المبيدات. وتوصي الدراسة الحالية باستخدام تجهيزه النيم لمكافحة خنفساء الدقيق الصدفية كبديل لاستخدام المبيدات الحشرية أو استخدامها ضمن برامج الإدارة المتكاملة للآفات.

تم اختبار فاعلية أربعة تجهيزات مختلفة من المبيدات المتوفرة تجارياً تحت الظروف المعملية ضد الحشرات الكاملة لخنفساء الدقيق الصدفية حيث تم إجراء التقييم الحيوي على درجة حرارة  $25 \pm 1$ °م و رطوبة نسبية قدرها  $70 \pm 5$ %, بواسطة خلط مبيد الفنفاليريت بمعدل 250 و 500 و 750 و 1000 مجم/لتر ، 25 و 50 و 75 و 100 مجم/لتر لمبيد الدايازينون ، 2500 و 5000 و 10000 و 20000 مجم/لتر للمبيد الميكروبي البيوفلاي ، 25 و 50 و 100 و 200 مجم/لتر للمبيد النيم ذو الأصل الطبيعي مع حبوب القمح . وقد تم تقدير النسبة المئوية للموت يومياً لمدة 10 أيام ثم حساب التركيز اللازم لقتل 50% و 95% وكذلك حساب الزمن اللازم لقتل 50% و 95% من يرقات خنفساء الدقيق الصدفية.

أوضحت النتائج أن النسبة المئوية للموت قد بلغت 100% بعد 10 أيام من المعاملة بتركيز قدره 1000 مجم/لتر من الفنفاليريت و بتركيز قدرها 50 و 75 و 100 مجم/لتر من الدايازينون. كما