

Evaluation of Chemical Control in A Rotational Program Against Whitefly on Tomato Crop

Farida A. Ayad¹, H.S. Taha¹, A. F.E. Afsah², A.R.I. Hanafy², S.A.G. Metwally² and Sherif Ayoub³

ABSTRACT

A chemical control program against *Bemisia Tabaci* was applied after treatment of tomato seedlings variety Rover F1 hybrid in the nursery using thiamethoxam and imidacloprid. In addition, six treatments with different rotational and conventional insecticides in a rotational program have been applied at Syngenta Kaha Station, Kalubia Governorate, Egypt. Two treatments were sprayed according to the proposed economic threshold (less than one adult per compound leaf) rotationally. Another two treatments were sprayed periodically every 6 days without taking into consideration the degree of infestation. The last two treatments were used as block application and sprayed every 6 days twice using the same chemical. The results showed that the best treatment was the block application, in the sequence of thiamethoxam, lambda-cyhalothrin, pymetrozine, pyriproxyfen and imidacloprid, fenpropathrin, pyriproxyfen, pymetrozine. They gave the highest efficacy on the adults of whitefly and obtained high yield of tomato crop. Regarding the virus infestation, both treatments showed the least number of infested plants with minimal surfaces of viral symptoms. The present study suggests that block application was a good rotational program in reducing the number of chemicals and reducing the hazard of pesticides to the environment.

INTRODUCTION

In recent years, the whitefly, *Bemisia tabaci* (Genn.) has become a serious pest of cotton, other field crops, vegetables, ornamentals and medical plants in Egypt. The high infestation results in significant reductions in total yield and quality. Furthermore, the adult stage acts as a vector of virus diseases for various hosts, and reductions in virus spread are usually achieved by vector control using insecticides (Cohen & Nitzani 1966; Gameel 1974; Shaheen, 1977 and Mazyad *et al.*, 1979 & 1986; Youssef *et al.*, 1979; El Nawawy *et al.*, 1979 and El Dakrouy *et al.*, 1983).

In most instances, however, the levels of vector control which are attainable with conventional insecticides are not sufficient to reduce virus spread to acceptable levels in either tomato nurseries where infestation first occurs, or in open production fields

(Sharaf and Allawi, 1981). At high summer temperature a generation of whiteflies can develop in only 2 to 3 weeks (Butler *et al.*, 1983). Also development of resistance to insecticides in whiteflies (Dittrich & Ernst 1983, Rowland *et al.*, 1991, Cahill *et al.*, 1995, Prabhaker *et al.*, 1998) has highlighted the need for an effective resistance management strategy (Prabhaker *et al.*, 1992). Integrated pest management (IPM) is the strongest enemy for resistance. Recently there is a great need for the wide use of IPM technique in pest control. This approach combines all the various tools and methods to manage pests at acceptable level. One of this tool is the development of a suitable insecticide rotation. First of all the spray program must begin when number of adults reached the economic thresholds (less than one / compound leaf). Therefore, the objective of this work is to compare the spray programs, according to the proposed economic threshold or every 6 days, with different classes of insecticides or as block application every 6 days twice using the same insecticide for whitefly control, virus reduction and on the yield of harvested tomatoes.

MATERIAL AND METHODS

An experiment was conducted on tomato variety Rover (E-446) F1 hybrid which can tolerate viral infestation. Seedlings were obtained from plastic houses at Fayoum Agricultural Research Center. Six treatments with different conventional insecticides in a rotational program were applied for the control of whitefly *Bemisia tabaci*. The field trial was conducted at Syngenta Kaha Station, Kalubia Governorate, Egypt.

Tomato seedlings were transplanted on open field plots on 23 rd August. The experimental design was strips of six treatments and two untreated check. Plot size was 375 m², four replicates were used and normal agricultural practices were followed. The rotational and conventional insecticides (Table 1) were applied in the season in a rotational program of eight treatments.

Treatments (A&B) were sprayed according to the proposed economic threshold (less than one adult / compound leaf)

¹Central Agricultural. Pesticides Laboratory Agric. Res. Center, Giza Egypt.

²Plant Protection Research Institute. Agric. Res. Center. Giza, Egypt.

³Syngenta Kaha Station, Kalubia Governorate, Egypt.

Project # FC/20/28-97, financed from Bureau de Liaison Agricole., Franco-Egyptien.

Received June 20, 2009, Accepted June 29, 2009

Table 1. Trade name, common name, formulation, chemical name and recommended rate of compounds used against *Bemisia tabaci* on tomato plantation

Trade name	Common name	Formulation	Chemical name	Recommended rate
Actelic	Pirimiphos-methyl	EC 50%	O-(2-(diethylamino)-6-methyl-4-pyrimidinyl) O,O-dimethyl phosphorothioate	375 ml / 100L water
Sumithion	Fenitrothion	EC 50%	O,O-dimethyl O-(3-methyl-4-nitrophenyl) phosphorothioate	1.5 L / feddan
Dathrin	Fenpropathrin	EC 20%		750 ml / feddan
Karate	Lambda-cyhalothrin	EC 5%	cyano(3-phenoxyphenyl)methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropane-1-carboxylate, (1A(S*),3A(Z))- (±)-	94 ml / 100 L water
Actara	Thiamethoxam	WG 25%	3-((2-chloro-5-thiazolyl)methyl)tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine	40 gm / 100 l
Confidor	Imidacloprid	SC 3.5%	1-(6-chloro-3-pyridinyl) methyl)-N-nitro-2-imidazolimidine	350 ml / feddan
Marshal	Carbosulfan	WP 50%	2,3-dihydro-2,2-dimethyl-7-benzofuranyl ((dib ty lamino)thio)methylcarbamate	150 gm/ 100 l
Admiral	Pyriproxyfen	EC 10%	2-(1-methyl-2-(4-phenoxyphenoxy)= ethoxy)pyridine	75 ml / 100 l water
Chess	Pymetrozine	WP 25%	4,5-dihydro-6-methyl-4-((3-pyridinyl= methylene)amino)-1,2,4-triazin-3(2H)-one, (E)-	480 gm / feddan
Polo	Diafenthiuron	SC 50%	N-(2,6-bis(1-methylethyl)-4-phenoxy= phenyl)-N-(1,1-dimethylethyl)-thiourea	300 ml / 100 l
Bemistop	Etofenprox		1-(2-(4-ethoxyphenyl)-2-methylpropoxy)= methyl)-3-phenoxybenzene	1 L / feddan
Evisect	Thiocyclam	SP50%	N,N-dimethyl-1,2,3-trithian-5-amine ethanediolate (1:1)	250 gm/100 L water
Lannate	Methomyl	WP 90%	methyl N-((methylamino)carbonyloxy)= ethanimidothioate	150 gm / 100 l water
Challenger	Chlorfenapyr	SC 36%	4-bromo-2-(4-chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)-1H-pyrrole-3-carbonitrile	
Match	Lufenuron	EC 5%	N-(((2,5-dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenyl)amino)= carbonyl)-(2,6-difluorobenzamide	200 ml / feddan

(C&D) were sprayed every 6 days .

(E&F) were sprayed as block application every 6 days using the same insecticide twice (during the generation of whitefly).

(G&H) were untreated [check], as illustrated in (Table 2).

Applications were done using a single nozzle knapsack sprayer with a spray volume of 200l / feddan.

The efficacy of the products used against the adult stage of whitefly was determined by counting insects on the lower surface of 20 compound leaves at the center of each replicate (80/treatment). Counts were made in the early morning when light activity is minimal according to Butler et al., (1988). Pre-treatment counts were made in the early morning just before application, and post treatment counts were made at three days intervals.

Percent reduction of adult stages of *Bemisia tabaci* was calculated for all treatment using the equation of Henderson & Tilton (1955). The virus symptoms were evaluated on 20 randomized plants in each replicate, at different time intervals 27,34,41,48,54, 63and 71 days after transplanting. Symptoms were evaluated morphologically. The number of plants exhibiting virus symptoms was recorded and percent plants showing virus symptoms estimates visually. The yield of the harvested tomatoes in each treatment was also recorded. The data of tomato yield were statistically analyzed according to Snedecore & Cochran (1980).

RESULTS AND DISCUSSION

Data in Table 3 showed the number of whitefly adults per 100 leaves and percent reduction of infestation at different time intervals. The best results were given by treatments E,F,D,B,C and A in descending order which mean that in the field the best treatment was the one of block application with 6 days intervals.

Treatment E was sprayed in the nursery with thiamethoxam (Actara) then sprayed in the field with twice applications of Actara, twice applications of Karate (lambda-cyhalothrin) twice applications of Chess (pymetrozine) and twice applications of Admiral (pyriproxyfen).the number of field applications was 8 during the season.

Treatment F was sprayed in the nursery with imidacloprid (Confidor). Then sprayed in the field twice applications of Confidor, with twice applications of Dathrin (fenpropathrin), twice applications of Admiral (pyriproxyfen) and twice applications of Chess (pymetrozine). Also, they number of field applications in this case was 8 during the season.

Treatment D was better than treatment B. Both treatments were treated with Confidor in the nursery and

in the field. Treatment D had only eight sprays while B had ten sprays (economic threshold). The last treatment C followed by A both were treated in the nursery with Actara and in the field treatment C had only eight sprays and treatment A had twelve sprays (economic threshold).

Regarding the rate of virus infection, in general these rates varied according to three factors: days after transplanting (incubation period), the insecticide used and the tomato variety. It is obvious from the average percentages of viral infected plants that treatment E was the least infected followed by B, D,C, A and F then the check.

The minimal severity of infection was obtained also by E and F followed by B, D, A, and C (equally) then the check (Table 4). Similar observation was found by Verma *et al.*, (1989), Rosset *et al.*(1990), and Dawood *et al.*,(1999). They reported that the incidence of tomato leaf curl virus was directly related to the population density of the vector *B. tabaci*

It is worthy to note that tomato plants resistant to TYLCV under field conditions is difficult to obtain Ahmed *et al.*, (1991) reported that 90% of the plants were highly susceptible, 9% were intermediate,0.5% were slightly susceptible and 0.3% showed no symptoms.

The effects of the tested programs on tomato yield were evaluated and presented in Table 5. The data showed that fruit yield of tomatoes were increased by all treatments. The highest yield was obtained after the application of program F which increased the yield by 174.17% (relative to untreated plots), while the lowest yield was recorded with program C which increased the yield by 81.83%. The data also showed no significant differences between the yield of A and D as well as between E and F. Generally it could be concluded that the programs applied on tomato are arranged according to the fruit yield (Kg) in the following order; F>E>D>A>B>C.

In conclusion, it can be deduced that treatment E gave the best results for whitefly control as well as highest yield (showing the least number of infested plants with minimal surface of virus symptoms) taking into consideration that it was treated in the field as block application sparing classes of insecticides, and reducing the number of sprays because they were well protected

Table 2. Rotation programs of the insecticides application

		Open field													
Application	Nursery 23.8	30.8	5.9	11.9	14.9	17.9	23.9	26.9	29.9	2.10	5.10	8.10	11.10	NO. of appl.	Programs followed
A	Actara	Karate	Chess	Admiral	Acetic Primphos- methyl	Confidor	Bemistop	Evisect	Polo	Lannate	Challenger	March	Marshal	12	Threshold
	Thiamethoxam	Lambda-cyhalothrin	Pymetrozine	Pyriproxyfen		Imidacloprid	etofenprox	Thio-cydam hydrogen oxalate	Diafenthiuron	Methomyl	Chlorfenapyr	Lufenuron	Carbosulfan		
B	Confidor	Datrin	Admiral	Chess		Sumithion	Actara	Evisect	Bemistop	Evisect	Polo	Lannate	Challenger	10	Threshold
	Imidacloprid	Fenpropathrin	Pyriproxyfen	Pymetrozine		Fenitrothion	Thiamethoxam		etofenprox	Thio-cydam hydrogen oxalate	Diafenthiuron	Methomyl	Chlorfenapyr		
C	Actara	Karate	Chess	Admiral	Acetic	Confidor	Confidor	Evisect	Bemistop	Evisect	Thio-cydam hydrogen oxalate		Polo	8	6 day intervals
	Thiamethoxam	Lambda-cyhalothrin	Pymetrozine	Pyriproxyfen		Imidacloprid	Imidacloprid		etofenprox	Thio-cydam hydrogen oxalate			Diafenthiuron		
D	Confidor	Datrin	Admiral	Chess		Sumithion	Actara	Evisect	Bemistop	Evisect	Thio-cydam hydrogen oxalate		Polo	8	6 day intervals
	Imidacloprid	Fenpropathrin	Pyriproxyfen	Pymetrozine		Fenitrothion	Thiamethoxam		etofenprox	Thio-cydam hydrogen oxalate			Diafenthiuron		
E	Actara	Karate	Chess	Admiral	Acetic	Confidor	Confidor	Evisect	Bemistop	Evisect	Thio-cydam hydrogen oxalate		Polo	8	Block appl. 6 day intervals
	Thiamethoxam	Lambda-cyhalothrin	Pymetrozine	Pyriproxyfen		Imidacloprid	Imidacloprid		etofenprox	Thio-cydam hydrogen oxalate			Diafenthiuron		
F	Confidor	Datrin	Admiral	Chess		Sumithion	Actara	Evisect	Bemistop	Evisect	Thio-cydam hydrogen oxalate		Polo	8	Block appl. 6 day intervals
	Imidacloprid	Fenpropathrin	Pyriproxyfen	Pymetrozine		Fenitrothion	Thiamethoxam		etofenprox	Thio-cydam hydrogen oxalate			Diafenthiuron		
G&H Check															

Table 3. Effect of insecticide rotation program on adults of whitefly

Treatment /100leaves	No of adult and		% Reduction																Average			
	0DDAI	3DDAI	6DDAI	9DDAI	12DDAI	15DDAI	18DDAI	21DDAI	24DDAI	27DDAI	30DDAI	33DDAI	36DDAI	39DDAI	42DDAI	45DDAI	48DDAI	51DDAI		54DDAI	57DDAI	60DDAI
A	No of adults	105	15	31.6	57	135	269	414	79	167	250	226	210	237	131	185	194	117	184			
	% Reduction		89	67	95	87	75	62	92	83	75	79	79	75	85	79	74	76	79			
B	No of adults	109	18	105	77	324	39	261	15	132	62	255	167	311	99	188	121	110	143			
	% Reduction		87	90	94	69	95	77	98	87	94	77	84	68	89	79	84	79	85			
C	No of adults	99	11	296	36	153	204	362	183	404	57	148	151	329	113	162	92	94	175			
	% Reduction		91	67	97	84	74	65	79	57	94	85	84	63	87	80	87	80	80			
D	No of adults	118	10	102	67	257	39	341	54	131	43	178	162	426	100	172	70	93	140			
	% Reduction		93	91	95	77	96	72	95	88	96	85	86	60	90	82	92	83	86			
E	No of adults	101	15	223	45	130	44	271	29	146	28	96	47	147	98	113	91	96	101			
	% Reduction		88	76	96	86	94	74	97	85	97	91	95	84	89	86	87	80	88			
F	No of adults	120	17	218	51	163	37	123	21	56	139	361	138	402	88	89	78	78	130			
	% Reduction		89	80	96	86	96	90	98	95	86	71	88	63	91	91	91	86	87			
G & H Check	No of adults	110	139	1010	1280	1049	865	1140	978	1045	1049	1127	1069	991	904	786	517	930				

Table 4. Effect of insecticides treatment in viral infection

Treatment	Total number of infected plants with virus												Percent surface area showing virus symptoms								
	27DAAI	34DAAI	41DAAI	48DAAI	54DAAI	63DAAI	71DAAI	77DAAI	84DAAI	91DAAI	98DAAI	104DAAI	112DAAI	120DAAI	127DAAI	134DAAI	141DAAI	148DAAI	155DAAI	162DAAI	
A	13	27	48	48	66	72	77	77	77	77	77	77	77	77	77	77	77	77	77	77	77
B	7	17	40	45	59	65	69	69	69	69	69	69	69	69	69	69	69	69	69	69	69
C	10	30	54	55	60	64	73	73	73	73	73	73	73	73	73	73	73	73	73	73	73
D	5	18	48	49	58	69	71	71	71	71	71	71	71	71	71	71	71	71	71	71	71
E	5	16	41	43	53	60	65	65	65	65	65	65	65	65	65	65	65	65	65	65	65
F	4	19	25	38	51	56	80	80	80	80	80	80	80	80	80	80	80	80	80	80	80
G & H Check	40	70	80	80	80	80	80	80	80	80	80	80	80	80	80	80	80	80	80	80	80

*DAAI= Days after Application

Table 5. Effect of insecticide treatments on tomato yield

Treatments	Harvested tomatoes			Yield in Kg	Yield %
	1st	2nd	3rd		
A	635	655	660	1950 ^b	122.86
B	588	579	570	1737 ^c	98.51
C	520	536	535	1591 ^d	81.83
D	651	668	658	1977 ^b	125.94
E	791	782	793	2366 ^a	170.40
F	800	801	798	2399 ^a	174.17
G & H Check	285	293	295	875 ^e	100

from the beginning with foliar sprays of neonicotinoids (only 8 sprays in the field during the season) these results may be helpful as an approach for managing insecticide resistance in *B.tabaci* population.

REFERENCES

- Ahmed, A.H.; M.S. Waif and Al-Izabi, F. (1991). Evaluation of wild domestic *lycopersicon* accessions for tomato yellow leaf curl virus resistance, Egypt, J. Hort. 18:23-43.
- Butler, G.DJr.; Henneberry, T.J. and Clayton, T.E. (1983). *Bemisia tabaci* (Homoptera: Aleyrodidae.) development, oviposition, and longevity in relation to temperature. Ann. Entomol. Soc. America., 76(2)310-313.
- Butler, G.D Jr.; Hutchinson, W.D; and Broza, M. (1988). Effects of aldicarb treatments to cotton on *Bemisia tabaci* and *Aphis gossypii* populations in Israel. Southwestern-Entomologist. 13: (2) 87-93.
- Cohen, S; and Nitzany, F.E. (1966). Transmission and host range of the tomato yellow leaf curl virus. Phytopathology 56, 1127-1131.
- Cahill, M.; Byrne, F.J.; Gorman, K.; Denholm, I.; and Devonshire, A.L. (1995). Pyrethroid and organophosphate resistance in the tobacco whitefly *Bemisia tabaci* (Homoptera: Aleyrodidae). Bulletin-of-Entomological-Research. 85: (2), 181-187.
- Dawood, M.Z.; K.K. El-Rafie; S.A. Aly And Hayder, M.F. (1999). Susceptibility of some tomato varieties and hybrids to whitefly *Bemisia tabaci* (Genn.) infestation in relation to rate of TYLCV infection and yield. Egypt J. Agric. Res. 77:1059-1064.
- Dittrich, V. and Ernst, G.H (1983). The resistance pattern in the whiteflies of Sudanese cotton on Mitt. Dtsch. Ges. Allg. Angew. Entomol.4:96-97.
- El- Nawawy, A.S.; O. Lamie, A. Salama, M.A. Ashry, E.A. Kadous, M. Darrag and M. Toulou (1979). Effect of several pesticides on the infestation with spider mites, whitefly, jassids and aphids in cotton fields in Kafr El-Shiekh Governorate. A.R. Egypt 1987. (Proc. 3rd Pesticide Conf., Tanta Univ., 1 :315-325).
- El- Dakrouy, F.; N.Z. Soliman and F.A. Khalil (1983). Effectiveness of certain insecticidal treatments against some pests attacking cotton plants. (Proc.5th Arab Pesticide Conf., Tanta Univ., 11:209-221).
- Gameel, O.I. (1974). Field evaluation of insecticide for jassid, *Empoasca lybica* De Berg, and whitefly, *Bemisia Tabaci* (Gennadius), control on cotton (Homoptera: Aleyrodidae). Bull. Ent. Soc. Egypt, Econ. Ser.,7:113-122.
- Henderson, C.F.; and Tilton, F.W. (1955). Tests with acaricides against the brown wheat mite. J. Econ. Entomol. 48:157-161.
- Mazyad, H.M. Omer, F., Al-Taher, K. and M. Salha (1979). Observation on the epidemiology of tomato yellow leaf curl diseases on tomato plants. Plant Dis. Repr., 63: 695-898.
- Mazyad, H.M., Nakhla, M.K. El-Amrety, A.A. and S.A. Doss (1986). Further studies on the epidemiology of tomato yellow leaf curl virus in Egypt. Acta Hort., 190:121-130.
- Prabhakar, N.; Toscano, N.C.; Perring, T.M.; Nuessey, G.; Kido, K. and Youngman, R.B. (1992). Resistance monitoring of the sweetpotato whitefly (Homoptera: Aleyrodidae) in the Imperial Valley of California. J. Econ. Entomol.; 85 (4). 1063-1068.
- Prabhakar, N.; Toscano, N.C.; and Henneberry, T.J. (1998). Evaluation of insecticide rotations and mixtures as resistance management strategies for *Bemisia argentifolii* (Hom: Aleyrodidae). J. Econ. Entomol. 91(4) 820-826.
- Rosset, P.; Eneses, R., Lastra, R., Gonzalez, W. (1990). Estimation of loss and identification of the geminivirus transmitted to tomato by the whitefly, *Bemisia tabaci* (Genn.) (Homoptera: Aleyrodidae) in Costa Rica. Manejo Ontegrado De Plagas, 15:24-35.
- Rowland, M.; Hackett, B. and Stribley, M. (1991). Evaluation of insecticides in field control simulators and standard laboratory bioassay against resistant and susceptible *Bemisia tabaci* (Homoptera : Aleyrodidae) from Sudan. Bulletin Entomological Res.81:2,189-199.
- Shaheen, A.H. (1977). Survey of pests attacking tomato in Egypt with some ecological notes. Agric. Res. Rev., Cairo, 55 (1): 49-57.
- Sharaf, N.S. and Allawi, T.F. (1980). Studies on whiteflies on tomato in the Jordan valley. 111 – laboratory and field experiments on the control of whitefly (*Bemisia tabaci* Genn, Homoptera: Aleyrodidae.) populations with organophosphorous insecticides and the incidence of the

- tomato yellow leaf curl virus. J.Pl. Dis.Prot.87(3):176-184.
- Snedecore,G.W. and Cochran (1980). "Statistical methods" 75th ed. Iowa State Univ., Press Ames, U.S.A.
- Youssef, K.H.; Hammad S.M. and Assem M.A. (1979). Studies on the control of the whitefly, *Bemisia tabaci* genn, on squash. (Proc.3rd Pesticide Conf., Tanta Univ., 1 : 370-376).
- Verma, A.K.; Basu, D.; Nath, P.S.; Ghatak S. and Mukhopadhyay S. (1989). Relationship between the population of whitefly, *Bemisia tabaci* (Genn.) (Homoptera: Aleyrodidae) and incidence of tomato leaf curl virus disease. Indian J. Mycological Res. 27:49-52.

الملخص العربي

تقييم المكافحة الكيميائية في برامج دوريه علي الذبابة البيضاء في محصول الطماطم

فريدة احمد عياد، حنان صلاح طه، عبد الجابر فتوح السيد عقصه، احمدحنفي، سامية جلال متولي، شريف ايوب

[C & D]. مبيدات تتبع مجاميع مختلفة.

بغض النظر عن نسبه الاصابه:

كان يتم رش النباتات كل ٦ أيام في تتابع دوري في المعاملتين

[E&F] كل رشتين كانتا بنفس المبيد في نفس جيل الحشرة block

application وتركت المعاملات [G&H] كمقارنه تم اخذ

متوسطها.

أوضحت النتائج أن أفضل المعاملات هي التي استخدمت

block application و قد أعطت أفضل نتيجة بالنسبة لمكافحه

الحشرات الكاملة كذلك المحصول مع انخفاض في شده الاصابه

الفيروسية. فاستخدام block application عمل علي خفض مجاميع

المبيدات المستخدمة ويعمل علي تأجيل تكون صفة المقاومة مع

تخفيض تأثير المبيدات علي البيئة.

تم تقييم المكافحة الكيميائية في برامج دوريه برش المبيدات

التقليدية وغير التقليدية في تتابع علي صنف الطماطم

Rover (E-446) F1 hybrid لمكافحه الذبابة البيضاء علي محصول

الطماطم في محطة بحوث Syngenta قها محافظه القلوبية. قسمت

المنطقة التجريبية إلي ثمانية معاملات، ثلاثة معاملات رشت في

المشتل بال Thiamethoxam وثلاثة معاملات رشت بالمشتل بال

Imidacloprid وذلك قبل نقلها إلي الأرض المستديمة ب ٤٨

ساعة، والمعاملتين الباقيتين تركنا كمقارنه بدون معاملة.

في الأرض المستديمة كان يتم رش النباتات في برنامج دوري

كلما اقترب عدد الحشرات الكاملة إلي الحد الاقتصادي الحرج

(متوسط حشره واحده / ورقه مركبه وذلك في معاملتين [A & B].

كان يتم رش النباتات كل ٦ أيام في برنامج دوري في المعاملتين