Date of Inspection	Mean no. individuals / plant (3 leaves)									General
	B. tabaci		T. urticae			Р.	Autographa	Т.	С.	Averages
	eggs	pupae	eggs	nymphs	adults	opercellula	spp.	tabaci	vulgaris	$(\overline{\mathbf{X}})$
Oct. 11,	6.7 a	20.7 ab	237.0 a	87.3 ab	58.0 b	3.0 a	0.3 a	0.0a	0.3 a	45.9 A
25,	3.7 ab	33.3 a	159.7 ab	142.0 a	137.7 a	0.0 a	1.0 a	0.3 a	0.0 a	53.1 A
Nov. 9,	0.7 b	16.7 b	109.0 ab	107.7 ab	33.7 bc	0.3 a	0.3 a	1.0 a	0.0 a	29.9 A
23,	5.7 a	2.7 b	3.3 b	2.3 b	3.0 c	0.0 a	0.0 a	0.0 a	0.0 a	1.9 B
Dec. 8,	6.0 a	4.7 b	0.7 b	1.7 b	0.3 c	0.0 a	0.0 a	0.0 a	0.0 a	1.5 B
22	0.7 b	5.0 b	0.0 b	1.7 b	0.0 c	0.0 a	0.0 a	0.0 a	0.0 a	0.8 B
$\overline{\mathbf{X}}$	8.9 D	13.9 CD	85.0 A	57.1 B	38.8 BC	0.6 D	0.3 D	0.2 D	0.1 D	22.2

Table 1. Pest species and predator with their densities on eggplant during autumn season, 2012.

Means followed by the same letter in each column are not significantly different at 0.05 level by LSD test.